

Analysis of the strong decays of the $Y(4660)$ in tetraquark scenario via the QCD sum rules*

Xiao-Song Yang (杨晓松)^{1,2} Zhi-Gang Wang (王志刚)[†]

¹Department of Physics, North China Electric Power University, Baoding 071003, China

²School of Nuclear Science and Engineering, North China Electric Power University, Beijing 102206, China

Abstract: Motivated by the enigmatic vector charmonium-like states, we investigate the strong decay behaviors of four types of vector tetraquark states, which are possible candidates for the $Y(4660)$, within the framework of three-point QCD sum rules based on rigorous quark-hadron duality. We take into account vacuum condensates up to dimension 5 on the QCD side and obtain the hadronic coupling constants and hence the partial decay widths of these states. The predicted total width, $61.5 \pm 7.3 \text{ MeV}$, is in excellent agreement with the experimental data for the $Y(4660)$, supporting its interpretation as a $[sc][\bar{s}\bar{c}]$ tetraquark state with $J^{PC} = 1^{--}$.

Keywords: tetraquark state, QCD sum rules

DOI: 10.1088/1674-1137/ae71a6 **CSTR:** 32044.14.ChinesePhysicsC.50093104

I. INTRODUCTION

In recent years, a number of charmonium-like states have been observed [1], which cannot be comfortably accommodated within the traditional quark model. They play an important role in understanding long-distance QCD dynamics and have inspired extensive studies, especially of the complicated Y states. In the present study, we focus on the $Y(4660)$ in the tetraquark scenario.

The $Y(4660)$ was first observed by the Belle collaboration in the process $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ in 2007 [2] and was later confirmed by the Belle, BaBar, and BESIII collaborations [3–5]. The masses and widths of the $Y(4660)$ and related states from different experiments are presented in Table 1.

In 2008, the Belle collaboration reported a measurement of the exclusive process $e^+e^- \rightarrow \Lambda_c^+\Lambda_c^-$ and observed a significant structure, which was denoted as $Y(4630)$ [6]. Owing to their similar masses and widths, the $Y(4630)$ and $Y(4660)$ are regarded as the same state by the Particle Data Group [1] and several groups [10–12]. However, the BESIII collaboration studied the process $e^+e^- \rightarrow \Lambda_c^+\bar{\Lambda}_c^-$ with higher statistics in 2023 [7], and the measured cross section indicated no enhancement around the $Y(4630)$ structure, which is significantly different from the Belle result [6].

In 2019, the Belle collaboration studied the $e^+e^- \rightarrow$

$D_s^+D_{s1}^-(2536)^-$ cross section and observed a charmonium-like state, $Y(4626)$, with a measured mass and width close to those of the $Y(4660)$ [8]. This was the first discovery of a Y state around 4.6 GeV in an open-charm channel. Later, the $Y(4626)$ was also confirmed in the $e^+e^- \rightarrow D_s^+D_{s2}^-(2573)^-$ channel [9]. If the $Y(4626)$ and $Y(4660)$ are the same state [1], it could be assigned as a $[sc][\bar{s}\bar{c}]$ state according to its decay into a $D_s^+D_{s1}^-$ pair.

On the theoretical side, after its discovery, the $Y(4660)$ was interpreted as a $\psi'f_0(980)$ molecular state [10, 13–15], a tetraquark state [11, 16–28], a hadro-charmonium state [29], $\psi(5S)$ [30–33], $\psi(6S)$ [34, 35], and so on.

In Ref. [36], the authors studied the mass spectrum of prospective hidden-bottom and hidden-charm hexaquark states via QCD sum rules and found that the $Y(4660)$ is close in magnitude to the $\Lambda_c\bar{\Lambda}_c$ -type baryonium state. The $\Lambda_c\bar{\Lambda}_c$ -type baryonium states have been studied in several works [37–40]. In Ref. [41], we also studied $\Lambda_c\bar{\Lambda}_c$ -type baryonium states via QCD sum rules and obtained a conclusion consistent with that of Ref. [36]. The interpretation of the $Y(4660)$ in the baryonium scenario provides meaningful insight into understanding the Y states.

In Table 2, we list the predictions for the masses of the $Y(4660)$ in the framework of QCD sum rules, which have achieved several successes in the study of exotic states [42–44]. It is clear that the experimental mass of

Received 1 May 2026; Accepted 19 May 2026; Accepted manuscript online 20 May 2026

* Supported by the National Natural Science Foundation of China (12575083)

[†] E-mail: zgwang@aliyun.com



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Article funded by SCOAP³ and published under licence by Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institute of Modern Physics of the Chinese Academy of Sciences and IOP Publishing Ltd

Table 1. The masses, widths, and channels measured in different experiments.

Year		Mass/MeV	Width/MeV	channel	Experiment
2007	$Y(4660)$	$4664 \pm 11 \pm 5$	$48 \pm 15 \pm 3$	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$	Belle[2]
2008	$Y(4630)$	4634^{+8+5}_{-7-8}	92^{+40+10}_{-24-21}	$e^+e^- \rightarrow \Lambda_c^+\Lambda_c^-$	Belle[6]
2023		not seen	not seen		BESIII[7]
2012	$Y(4660)$	$4669 \pm 21 \pm 3$	$104 \pm 48 \pm 10$	$e^+e^- \rightarrow \psi(2S)\pi^+\pi^-$	BaBar[3]
2014	$Y(4660)$	$4652 \pm 10 \pm 8$	$68 \pm 11 \pm 1$	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$	Belle[4]
2019	$Y(4626)$	$4625.9^{+6.2}_{-6.0} \pm 0.4$	$49.8^{+13.9}_{-11.5} \pm 4.0$	$e^+e^- \rightarrow D_s^+D_{s1}(2536)^-$	Belle [8]
2020	$Y(4626)$	$4619.8^{+8.9}_{-8.0} \pm 2.3$	$47.0^{+31.3}_{-14.8} \pm 4.6$	$e^+e^- \rightarrow D_s^+D_{s2}^*(2573)^-$	Belle [9]
2021	$Y(4660)$	$4651.0 \pm 37.8 \pm 2.1$	$155.4 \pm 24.8 \pm 0.8$	$e^+e^- \rightarrow \pi^+\pi^-\psi(3686)$	BESIII
2023	$Y(4660)$	4675.3 ± 29.5	218.3 ± 72.9	$e^+e^- \rightarrow D^{*0}D^{*-}\pi^+$	BESIII[5]

Table 2. The masses are obtained from the QCD sum rules with different quark structures, where OPE denotes the truncation of the operator product expansion up to vacuum condensates of dimension n , and No indicates that the vacuum condensates of dimension n' are not included.

	Structures	OPE (No)	mass/GeV	References
$Y(4660)$	$\psi' f_0(980)$	10	4.71	[15]
		6	4.67	[14]
$Y(4660)$	$[sc]_S[\bar{s}\bar{c}]_V + [sc]_V[\bar{s}\bar{c}]_S$	8 (7)	4.65	[16]
		10	4.68	[21]
$Y(4660)$	$[sc]_{\bar{V}}[\bar{s}\bar{c}]_A - [sc]_A[\bar{s}\bar{c}]_{\bar{V}}$	8 (7)	4.64	[18]
$Y(4660)$	$[qc]_S[\bar{q}\bar{c}]_V + [qc]_V[\bar{q}\bar{c}]_S$	8 (7)	4.64	[18]
$Y(4360)$	$[qc]_S[\bar{q}\bar{c}]_V + [qc]_V[\bar{q}\bar{c}]_S$	10	4.34	[23]
$Y(4660)$	$[sc]_P[\bar{s}\bar{c}]_A - [sc]_A[\bar{s}\bar{c}]_P$	10	4.70	[24]
			4.66	[23]
			4.66	[24]
$Y(4660)$	$[qc]_P[\bar{q}\bar{c}]_A - [qc]_A[\bar{q}\bar{c}]_P$	10	4.59	[23]
			4.66	[26]
$Y(4660)$	$[qc]_A[\bar{q}\bar{c}]_A$	10	4.66	[25]
			4.69	[26]
$Y(4660)$	$[sc]_S[\bar{s}\bar{c}]_S$	6	4.69	[19]
$Y(4660)$	$[sc]_{\bar{A}}[\bar{s}\bar{c}]_V + [sc]_V[\bar{s}\bar{c}]_{\bar{A}}$	10	4.65	[27]
$Y(4660)$	$[sc]_S[\bar{s}\bar{c}]_{\bar{V}} - [sc]_{\bar{V}}[\bar{s}\bar{c}]_S$	10	4.68	[27]
$Y(4660)$	$\Lambda_c\bar{\Lambda}_c$	12	4.78	[36]
$Y(4660)$	$\Lambda_c\bar{\Lambda}_c$	16	4.68	[41]

the $Y(4660)$ can be reproduced with different structures; thus, we should explore its decay widths to shed light on its nature. Alternatively, the $Y(4660)$ might have several important Fock components, such as molecular and tetraquark components, and embody their collective effects.

Considering that the $Y(4260)$ and $Y(4660)$ decay into $J/\psi\pi^+\pi^-$ and $\psi(2S)\pi^+\pi^-$, respectively [2, 45], one might assign the $Y(4660)$ as the radial excitation of the $Y(4260)$ [22]. However, the process $Y(4230) \rightarrow \psi(2S)\pi^+\pi^-$ was observed by the BESIII collaboration [46–47]. In Ref. [48],

we investigated the ground states and first radial excitations of the vector hidden-charm tetraquark states with an explicit P -wave via QCD sum rules. The results support assigning the $Y(4260)$ as the $1P$ tetraquark state, leaving no room to accommodate the $Y(4660)$ among the first radial excitations.

In Refs. [23–25, 48–50], we studied the mass spectrum of the vector hidden-charm tetraquark states with or without an explicit P -wave via QCD sum rules, and the results support assigning the $Y(4660)$ as the $[sc]_P[\bar{s}\bar{c}]_A$ –

$[sc]_A[\bar{s}\bar{c}]_P$ type tetraquark state. In Ref. [51], we studied the decay behaviors of the $[sc]_P[\bar{s}\bar{c}]_A - [sc]_A[\bar{s}\bar{c}]_P$ type tetraquark state. Later, in Refs. [26–27], we took the scalar (S), pseudoscalar (P), vector (V), axial-vector (A), and tensor ($\tilde{A}, \tilde{V}$) (anti)diquark operators as the elementary building blocks to construct local four-quark currents with or without hidden strangeness in a comprehensive and consistent way, and updated the previous calculations. The $\tilde{A}$ and $\tilde{V}$ denote the $J^P = 1^+$ and $J^P = 1^-$ components of the tensor diquarks $\varepsilon^{ijk}q_j^T C\sigma_{\mu\nu}c_k$ or $\varepsilon^{ijk}q_j^T C\sigma_{\mu\nu}\gamma_5 c_k$. We studied the mass spectrum of those tetraquark states with $J^{PC} = 1^{--}$ and 1^{-+} in detail and revisited the assignments of the Y states. The results favor assigning the $Y(4660)$ as the $[sc]_{\tilde{A}}[\bar{s}\bar{c}]_V + [sc]_V[\bar{s}\bar{c}]_{\tilde{A}}$, $[sc]_S[\bar{s}\bar{c}]_{\tilde{V}} - [sc]_{\tilde{V}}[\bar{s}\bar{c}]_S$, $[uc]_P[\bar{u}\bar{c}]_A + [dc]_P[\bar{d}\bar{c}]_A - [uc]_A[\bar{u}\bar{c}]_P - [dc]_A[\bar{d}\bar{c}]_P$, or $[uc]_A[\bar{u}\bar{c}]_A + [dc]_A[\bar{d}\bar{c}]_A$ type tetraquark state.

In the present work, we take these four tetraquark configurations into account to explore the strong decay behaviors of the $Y(4660)$. We study the hadronic coupling constants and partial decay widths of the two-body

strong decays, which can occur through the Okubo-Zweig-Iizuka super-allowed fall-apart mechanism without annihilation or creation of a quark-antiquark pair. Specifically, we take into account the channels $Y \rightarrow \bar{D}_{(s)}D_{(s)}, \bar{D}_{(s)}^*D_{(s)}, \bar{D}_{(s)}^*D_{(s)}^*, \bar{D}_{(s)0}D_{(s)}^*, \bar{D}_{(s)1}D_{(s)}, \eta_c\omega(\phi(1020)), J/\psi\omega(\phi(1020)), \chi_{c0}\omega(\phi(1020)), \chi_{c1}\omega(\phi(1020))$, and $J/\psi f_0(500)(f_0(980))$.

The article is organized as follows: in Section II, we obtain the hadronic coupling constants in the two-body strong decays of the four vector tetraquark states via QCD sum rules; in Section III, we present the numerical results and discussion; finally, the conclusion is presented in Section IV.

II. QCD SUM RULES FOR THE HADRONIC COUPLING CONSTANTS

We choose the following four tetraquark currents with quantum numbers $J^{PC} = 1^{--}$ to study the $Y(4660)$:

$$J_\mu^{PA}(x) = \frac{\varepsilon^{ijk}\varepsilon^{lmn}}{2} \left[u_j^T(x) C c_k(x) \bar{u}_m(x) \gamma_\mu C \bar{c}_n^T(x) + d_j^T(x) C c_k(x) \bar{d}_m(x) \gamma_\mu C \bar{c}_n^T(x) - u_j^T(x) C \gamma_\mu c_k(x) \bar{u}_m(x) C \bar{c}_n^T(x) - d_j^T(x) C \gamma_\mu c_k(x) \bar{d}_m(x) C \bar{c}_n^T(x) \right], \quad (1)$$

$$J_\mu^{AA}(x) = \frac{\varepsilon^{ijk}\varepsilon^{lmn}}{2} \left[u_j^T(x) C \gamma_\mu c_k(x) \bar{u}_m(x) \gamma_\nu C \bar{c}_n^T(x) + d_j^T(x) C \gamma_\mu c_k(x) \bar{d}_m(x) \gamma_\nu C \bar{c}_n^T(x) - u_j^T(x) C \gamma_\nu c_k(x) \bar{u}_m(x) \gamma_\mu C \bar{c}_n^T(x) - d_j^T(x) C \gamma_\nu c_k(x) \bar{d}_m(x) \gamma_\mu C \bar{c}_n^T(x) \right], \quad (2)$$

$$J_\mu^{\tilde{A}V}(x) = \frac{\varepsilon^{ijk}\varepsilon^{lmn}}{\sqrt{2}} \left[s_j^T(x) C \sigma_{\mu\nu} \gamma_5 c_k(x) \bar{s}_m(x) \gamma_5 \gamma^\nu C \bar{c}_n^T(x) + s_j^T(x) C \gamma^\nu \gamma_5 c_k(x) \bar{s}_m(x) \gamma_5 \sigma_{\mu\nu} C \bar{c}_n^T(x) \right], \quad (3)$$

$$J_\mu^{\tilde{S}\tilde{V}}(x) = \frac{\varepsilon^{ijk}\varepsilon^{lmn}}{\sqrt{2}} \left[s_j^T(x) C \gamma_5 c_k(x) \bar{s}_m(x) \sigma_{\mu\nu} C \bar{c}_n^T(x) - s_j^T(x) C \sigma_{\mu\nu} c_k(x) \bar{s}_m(x) \gamma_5 C \bar{c}_n^T(x) \right], \quad (4)$$

where i, j, k, m and n are color indices, and the superscripts S, P, A ($\tilde{A}$), and V ($\tilde{V}$) denote scalar, pseudoscalar, axialvector, and vector diquarks or antidiquarks, respectively. For conventional mesons, we adopt the following currents,

$$\begin{aligned} J^{\tilde{D}}(x) &= \bar{c}(x) i \gamma_5 u(x), \\ J^D(y) &= \bar{u}(y) i \gamma_5 c(y), \\ J_\alpha^{\tilde{D}^*}(x) &= \bar{c}(x) \gamma_\alpha u(x), \\ J_\beta^{D^*}(y) &= \bar{u}(y) \gamma_\beta c(y), \end{aligned}$$

$$\begin{aligned} J^{\tilde{D}_0}(x) &= \bar{c}(x) u(x), \\ J_\alpha^{\tilde{D}_1}(x) &= \bar{c}(x) \gamma_\alpha \gamma_5 u(x), \\ J^{\tilde{D}_s}(x) &= \bar{c}(x) i \gamma_5 s(x), \\ J^{D_s}(y) &= \bar{s}(y) i \gamma_5 c(y), \\ J_\alpha^{\tilde{D}_s^*}(x) &= \bar{c}(x) \gamma_\alpha s(x), \\ J_\beta^{D_s^*}(y) &= \bar{s}(y) \gamma_\beta c(y), \\ J^{\tilde{D}_{s0}}(x) &= \bar{c}(x) s(x), \\ J_\alpha^{\tilde{D}_{s1}}(x) &= \bar{c}(x) \gamma_\alpha \gamma_5 s(x), \end{aligned}$$

$$\begin{aligned}
J^{\eta_c}(x) &= \bar{c}(x) i \gamma_5 c(x), \\
J_\alpha^{J/\psi}(x) &= \bar{c}(x) \gamma_\alpha c(x), \\
J^{\chi_{c0}}(x) &= \bar{c}(x) c(x), \\
J_\alpha^{\chi_{c1}}(x) &= \bar{c}(x) \gamma_\alpha \gamma_5 c(x), \\
J_\alpha^\omega(y) &= \frac{\bar{u}(y) \gamma_\alpha u(y) + \bar{d}(y) \gamma_\alpha d(y)}{\sqrt{2}}, \\
J_\alpha^{\phi(1020)}(y) &= \bar{s}(y) \gamma_\alpha s(y), \\
J_{f_0(500)}(y) &= \frac{\bar{u}(y) u(y) + \bar{d}(y) d(y)}{\sqrt{2}}, \\
J_{f_0(980)}(y) &= \bar{s}(y) s(y).
\end{aligned} \tag{5}$$

Now we introduce the following three-point correlation functions to study the hadronic coupling constants within QCD sum rules,

$$\Pi_\mu^{DDPA}(p, q) = i^2 \int d^4x d^4y e^{ip \cdot x} e^{iq \cdot y} \langle 0 | T \{ J^D(x) J^D(y) J_\mu^{PA\dagger}(0) \} | 0 \rangle, \tag{6}$$

$$\Pi_{\alpha\mu}^{\bar{D}^* DPA}(p, q) = i^2 \int d^4x d^4y e^{ip \cdot x} e^{iq \cdot y} \langle 0 | T \{ J_\alpha^{\bar{D}^*}(x) J^D(y) J_\mu^{PA\dagger}(0) \} | 0 \rangle, \tag{7}$$

$$\Pi_{\alpha\beta\mu}^{\bar{D}^* D^* PA}(p, q) = i^2 \int d^4x d^4y e^{ip \cdot x} e^{iq \cdot y} \langle 0 | T \{ J_\alpha^{\bar{D}^*}(x) J_\beta^{D^*}(y) J_\mu^{PA\dagger}(0) \} | 0 \rangle, \tag{8}$$

$$\Pi_{\alpha\mu}^{\bar{D}_0 D^* PA}(p, q) = i^2 \int d^4x d^4y e^{ip \cdot x} e^{iq \cdot y} \langle 0 | T \{ J_\alpha^{\bar{D}_0}(x) J_\mu^{D^*}(y) J_\mu^{PA\dagger}(0) \} | 0 \rangle, \tag{9}$$

$$\Pi_{\alpha\mu}^{\bar{D}_1 DPA}(p, q) = i^2 \int d^4x d^4y e^{ip \cdot x} e^{iq \cdot y} \langle 0 | T \{ J_\alpha^{\bar{D}_1}(x) J^D(y) J_\mu^{PA\dagger}(0) \} | 0 \rangle, \tag{10}$$

$$\Pi_{\alpha\mu}^{\eta_c \omega PA}(p, q) = i^2 \int d^4x d^4y e^{ip \cdot x} e^{iq \cdot y} \langle 0 | T \{ J^{\eta_c}(x) J_\alpha^\omega(y) J_\mu^{PA\dagger}(0) \} | 0 \rangle, \tag{11}$$

$$\Pi_{\alpha\beta\mu}^{J/\psi \omega PA}(p, q) = i^2 \int d^4x d^4y e^{ip \cdot x} e^{iq \cdot y} \langle 0 | T \{ J_\alpha^{J/\psi}(x) J_\beta^\omega(y) J_\mu^{PA\dagger}(0) \} | 0 \rangle, \tag{12}$$

$$\Pi_{\alpha\mu}^{\chi_{c0} \omega PA}(p, q) = i^2 \int d^4x d^4y e^{ip \cdot x} e^{iq \cdot y} \langle 0 | T \{ J^{\chi_{c0}}(x) J_\alpha^\omega(y) J_\mu^{PA\dagger}(0) \} | 0 \rangle, \tag{13}$$

$$\Pi_{\alpha\beta\mu}^{\chi_{c1} \omega PA}(p, q) = i^2 \int d^4x d^4y e^{ip \cdot x} e^{iq \cdot y} \langle 0 | T \{ J_\alpha^{\chi_{c1}}(x) J_\beta^\omega(y) J_\mu^{PA\dagger}(0) \} | 0 \rangle, \tag{14}$$

$$\Pi_{\alpha\mu}^{J/\psi f_0(500) PA}(p, q) = i^2 \int d^4x d^4y e^{ip \cdot x} e^{iq \cdot y} \langle 0 | T \{ J_\alpha^{J/\psi}(x) J_{f_0(500)}(y) J_\mu^{PA\dagger}(0) \} | 0 \rangle. \tag{15}$$

We can obtain the other correlation functions by making the following replacements: $\mu \rightarrow \mu\nu$, $PA \rightarrow AA$ for the current $J_{\mu\nu}^{AA}$; ($\bar{D}$, D , $\bar{D}^*$, D^* , $\bar{D}_0$, $\bar{D}_1$, ω , $f_0(500)$) $\rightarrow$ ($\bar{D}_s$, D_s , $\bar{D}_s^*$, D_s^* , $\bar{D}_{s0}$, $\bar{D}_{s1}$, $\phi(1020)$, $f_0(980)$), $PA \rightarrow \bar{A}V$ for the current J_μ^{AV} ; and $\mu \rightarrow \mu\nu$, ($\bar{D}$, D , $\bar{D}^*$, D^* , $\bar{D}_0$, $\bar{D}_1$, ω , $f_0(500)$) $\rightarrow$ ($\bar{D}_s$, D_s , $\bar{D}_s^*$, D_s^* , $\bar{D}_{s0}$, $\bar{D}_{s1}$, $\phi(1020)$, $f_0(980)$), $PA \rightarrow \bar{S} \tilde{V}$ for the current $J_{\mu\nu}^{S\tilde{V}}$.

On the phenomenological side, we insert a complete set of intermediate hadronic states with the same quantum

numbers as the interpolating currents and adopt the following definitions of the decay constants or pole residues,

$$\begin{aligned}
\langle 0 | J^S(0) | S(p) \rangle &= f_S m_S, \\
\langle 0 | J^P(0) | P(p) \rangle &= \frac{f_P m_P^2}{m_c}, \\
\langle 0 | J_\alpha^A(0) | A(p) \rangle &= f_A m_A \xi_\alpha, \\
\langle 0 | J_\alpha^V(0) | V(p) \rangle &= f_V m_V \xi_\alpha,
\end{aligned} \tag{16}$$

$$\begin{aligned}
\langle 0 | J_{\mu}^{PA/\tilde{A}V}(0) | Y_{PA/\tilde{A}V}(p') \rangle &= \lambda_{PA/\tilde{A}V} \varepsilon_{\mu}, \\
\langle 0 | J_{\mu\nu}^{S\tilde{V}}(0) | Y_{S\tilde{V}}(p') \rangle &= \lambda_{S\tilde{V}} \varepsilon_{\mu\nu\alpha\beta} \varepsilon^{\alpha} p'^{\beta}, \\
\langle 0 | J_{\mu\nu}^{S\tilde{V}}(0) | X_{S\tilde{A}}(p') \rangle &= \tilde{\lambda}_{S\tilde{A}} (\varepsilon_{\mu} p'_{\nu} - \varepsilon_{\nu} p'_{\mu}), \\
\langle 0 | J_{\mu\nu}^{AA}(0) | Y_{AA}(p') \rangle &= \lambda_{AA} (\varepsilon_{\mu} p'_{\nu} - \varepsilon_{\nu} p'_{\mu}), \\
\langle 0 | J_{\mu\nu}^{AA}(0) | X_{AA}(p') \rangle &= \tilde{\lambda}_{AA} \varepsilon_{\mu\nu\alpha\beta} \varepsilon^{\alpha} p'^{\beta}, \quad (17)
\end{aligned}$$

where ξ_{μ} and ε_{μ} denote the polarization vectors of the corresponding mesons or tetraquark states; S denotes the scalar mesons, including D_0 , D_{s0} , χ_{c0} , $f_0(500)$, and $f_0(980)$; P denotes the pseudoscalar mesons, including D , D_s , and η_c (note that we should make the replacement $m_c \rightarrow m_c + m_s$ for the D_s meson and $m_c \rightarrow 2m_c$ for the η_c meson); A denotes the axial-vector mesons, including D_1 , D_{s1} , and χ_{c1} ; and V denotes the vector mesons, including D^* , D_s^* , J/ψ , ω , and $\phi(1020)$. The X_{AA} and $X_{S\tilde{A}}$ denote tetraquark states with $J^{PC} = 1^{+-}$, which can couple to the currents $J_{\mu\nu}^{AA}$ and $J_{\mu\nu}^{S\tilde{V}}$, respectively. Furthermore, the hadronic coupling constants are defined as follows,

$$\begin{aligned}
\langle \bar{D}(p) D(q) | Y_{PA}(p') \rangle &= i(p-q) \cdot \varepsilon G_{\bar{D}DPA}, \\
\langle \bar{D}(p) D(q) | Y_{AA}(p') \rangle &= -i(p-q) \cdot \varepsilon G_{\bar{D}DAA}, \\
\langle \bar{D}_s(p) D_s(q) | Y_{\tilde{A}V}(p') \rangle &= (p-q) \cdot \varepsilon G_{\bar{D}_s D_s \tilde{A}V}, \\
\langle \bar{D}_s(p) D_s(q) | Y_{S\tilde{V}}(p') \rangle &= i(p-q) \cdot \varepsilon G_{\bar{D}_s D_s S\tilde{V}}, \quad (18)
\end{aligned}$$

$$\begin{aligned}
\langle \bar{D}^*(p) D(q) | Y_{PA}(p') \rangle &= -\varepsilon^{\lambda\tau\rho\sigma} p_{\lambda} \xi_{\tau}^* p'_{\rho} \varepsilon_{\sigma} G_{\bar{D}^* DPA}, \\
\langle \bar{D}^*(p) D(q) | Y_{AA}(p') \rangle &= -i\varepsilon^{\lambda\tau\rho\sigma} p_{\lambda} \xi_{\tau}^* p'_{\rho} \varepsilon_{\sigma} G_{\bar{D}^* DAA}, \\
\langle \bar{D}_s^*(p) D_s(q) | Y_{\tilde{A}V}(p') \rangle &= -\varepsilon^{\lambda\tau\rho\sigma} p_{\lambda} \xi_{\tau}^* p'_{\rho} \varepsilon_{\sigma} G_{\bar{D}_s^* D_s \tilde{A}V}, \\
\langle \bar{D}_s^*(p) D_s(q) | Y_{S\tilde{V}}(p') \rangle &= -i\varepsilon^{\lambda\tau\rho\sigma} p_{\lambda} \xi_{\tau}^* p'_{\rho} \varepsilon_{\sigma} G_{\bar{D}_s^* D_s S\tilde{V}}. \quad (19)
\end{aligned}$$

For simplicity, the definitions of the other hadronic coupling constants are given in Appendix A.

By isolating the ground-state contributions, we obtain the following correlation functions [52–53]:

$$\Pi_{\mu}^{\bar{D}DPA}(p, q) = \Pi_{\bar{D}DPA}(p'^2, p^2, q^2) (q-p)_{\mu} + \dots, \quad (20)$$

$$\Pi_{\alpha\mu}^{\bar{D}^*DPA}(p, q) = \Pi_{\bar{D}^*DPA}(p'^2, p^2, q^2) (-i\varepsilon_{\alpha\mu\lambda\tau} p^{\lambda} q^{\tau}) + \dots, \quad (21)$$

$$\Pi_{\alpha\beta\mu}^{\bar{D}^*D^*PA}(p, q) = \Pi_{\bar{D}^*D^*PA}(p'^2, p^2, q^2) (-g_{\alpha\beta} p_{\mu}) + \dots, \quad (22)$$

$$\Pi_{\alpha\mu}^{\bar{D}_0D^*PA}(p, q) = \Pi_{\bar{D}_0D^*PA}(p'^2, p^2, q^2) (-ig_{\alpha\mu} p \cdot q) + \dots, \quad (23)$$

$$\Pi_{\alpha\mu}^{\bar{D}_1DPA}(p, q) = \Pi_{\bar{D}_1DPA}(p'^2, p^2, q^2) (ig_{\alpha\mu}) + \dots, \quad (24)$$

$$\Pi_{\alpha\mu}^{\eta_c\omega PA}(p, q) = \Pi_{\eta_c\omega PA}(p'^2, p^2, q^2) (\varepsilon_{\alpha\mu\lambda\tau} p^{\lambda} q^{\tau}) + \dots, \quad (25)$$

$$\Pi_{\alpha\beta\mu}^{J/\psi\omega PA}(p, q) = \Pi_{J/\psi\omega PA}(p'^2, p^2, q^2) (ig_{\alpha\beta} p_{\mu}) + \dots, \quad (26)$$

$$\Pi_{\alpha\mu}^{\chi_{c0}\omega PA}(p, q) = \Pi_{\chi_{c0}\omega PA}(p'^2, p^2, q^2) (g_{\alpha\mu}) + \dots, \quad (27)$$

$$\Pi_{\alpha\beta\mu}^{\chi_{c1}\omega PA}(p, q) = \Pi_{\chi_{c1}\omega PA}(p'^2, p^2, q^2) (-i\varepsilon_{\alpha\beta\mu\lambda} p^{\lambda} p \cdot q) + \dots, \quad (28)$$

$$\Pi_{\alpha\mu}^{J/\psi f_0(500)PA}(p, q) = \Pi_{J/\psi f_0(500)PA}(p'^2, p^2, q^2) (-g_{\alpha\mu}) + \dots. \quad (29)$$

The other correlation functions for the currents $J_{\mu\nu}^{AA}$, $J_{\mu}^{\tilde{A}V}$, and $J_{\mu\nu}^{S\tilde{V}}$ are presented in Appendix B. In the above equations, the scalar invariant components are expressed as follows:

$$\Pi_{\bar{D}DPA}(p'^2, p^2, q^2) = \frac{\lambda_{\bar{D}DPA}}{(m_Y^2 - p'^2)(m_D^2 - p^2)(m_D^2 - q^2)} + \frac{C_{\bar{D}DPA}}{(m_D^2 - p^2)(m_D^2 - q^2)} + \dots, \quad (30)$$

$$\Pi_{\bar{D}^*DPA}(p'^2, p^2, q^2) = \frac{\lambda_{\bar{D}^*DPA}}{(m_Y^2 - p'^2)(m_{D^*}^2 - p^2)(m_D^2 - q^2)} + \frac{C_{\bar{D}^*DPA}}{(m_{D^*}^2 - p^2)(m_D^2 - q^2)} + \dots, \quad (31)$$

$$\Pi_{\bar{D}^*D^*PA}(p'^2, p^2, q^2) = \frac{\lambda_{\bar{D}^*D^*PA}}{(m_Y^2 - p'^2)(m_{D^*}^2 - p^2)(m_{D^*}^2 - q^2)} + \frac{C_{\bar{D}^*D^*PA}}{(m_{D^*}^2 - p^2)(m_{D^*}^2 - q^2)} + \dots, \quad (32)$$

$$\Pi_{\bar{D}_0D^*PA}(p'^2, p^2, q^2) = \frac{\lambda_{\bar{D}_0D^*PA}}{(m_Y^2 - p'^2)(m_{D_0}^2 - p^2)(m_{D^*}^2 - q^2)} + \frac{C_{\bar{D}_0D^*PA}}{(m_{D_0}^2 - p^2)(m_{D^*}^2 - q^2)} + \dots, \quad (33)$$

$$\Pi_{\bar{D}_1 DPA}(p'^2, p^2, q^2) = \frac{\lambda_{\bar{D}_1 DPA}}{(m_Y^2 - p'^2)(m_{D_1}^2 - p^2)(m_D^2 - q^2)} + \frac{C_{\bar{D}_1 DPA}}{(m_{D_1}^2 - p^2)(m_D^2 - q^2)} + \dots, \quad (34)$$

$$\Pi_{\eta_c \omega PA}(p'^2, p^2, q^2) = \frac{\lambda_{\eta_c \omega PA}}{(m_Y^2 - p'^2)(m_{\eta_c}^2 - p^2)(m_\omega^2 - q^2)} + \frac{C_{\eta_c \omega PA}}{(m_{\eta_c}^2 - p^2)(m_\omega^2 - q^2)} + \dots, \quad (35)$$

$$\Pi_{J/\psi \omega PA}(p'^2, p^2, q^2) = \frac{\lambda_{J/\psi \omega PA}}{(m_Y^2 - p'^2)(m_{J/\psi}^2 - p^2)(m_\omega^2 - q^2)} + \frac{C_{J/\psi \omega PA}}{(m_{J/\psi}^2 - p^2)(m_\omega^2 - q^2)} + \dots, \quad (36)$$

$$\Pi_{\chi_{c0} \omega PA}(p'^2, p^2, q^2) = \frac{\lambda_{\chi_{c0} \omega PA}}{(m_Y^2 - p'^2)(m_{\chi_{c0}}^2 - p^2)(m_\omega^2 - q^2)} + \frac{C_{\chi_{c0} \omega PA}}{(m_{\chi_{c0}}^2 - p^2)(m_\omega^2 - q^2)} + \dots, \quad (37)$$

$$\Pi_{\chi_{c1} \omega PA}(p'^2, p^2, q^2) = \frac{\lambda_{\chi_{c1} \omega PA}}{(m_Y^2 - p'^2)(m_{\chi_{c1}}^2 - p^2)(m_\omega^2 - q^2)} + \frac{C_{\chi_{c1} \omega PA}}{(m_{\chi_{c1}}^2 - p^2)(m_\omega^2 - q^2)} + \dots, \quad (38)$$

$$\Pi_{J/\psi f_0(500) PA}(p'^2, p^2, q^2) = \frac{\lambda_{J/\psi f_0(500) PA}}{(m_Y^2 - p'^2)(m_{J/\psi}^2 - p^2)(m_{f_0(500)}^2 - q^2)} + \frac{C_{J/\psi f_0(500) PA}}{(m_{J/\psi}^2 - p^2)(m_{f_0(500)}^2 - q^2)} + \dots. \quad (39)$$

By making the replacement $PA \rightarrow AA$, we can obtain the corresponding scalar invariant components for the current $J_{\mu\nu}^{AA}$. With the replacements $(\bar{D}, D, \bar{D}^*, D^*, \bar{D}_0, \bar{D}_1, \omega, f_0(500)) \rightarrow (\bar{D}_s, D_s, \bar{D}_s^*, D_s^*, \bar{D}_{s0}, \bar{D}_{s1}, \phi(1020), f_0(980))$ and $PA \rightarrow \tilde{A}\tilde{V}(S\tilde{V})$, we can obtain the corresponding scalar invariant components for the current $J_{\mu\nu}^{A\tilde{V}}(J_{\mu\nu}^{S\tilde{V}})$.

Since both currents $J_{\mu\nu}^{AA}$ and $J_{\mu\nu}^{S\tilde{V}}$ can potentially couple to the tetraquark states with the quantum numbers $J^{PC} = 1^{+-}$ and $J^{PC} = 1^{+-}$, we cannot clearly eliminate the contamination from the axial-vector components in the $Y_{AA} \rightarrow \bar{D}_0 D^*$, $Y_{AA} \rightarrow \bar{D}_1 D$, $Y_{AA} \rightarrow \chi_{c0} \omega$, $Y_{AA} \rightarrow J/\psi f_0(500)$ and $Y_{S\tilde{V}} \rightarrow \eta_c \phi$ channels in the calculations. Therefore, we parameterize the contributions of the axial-vector com-

ponents on the hadron side as $\bar{\lambda}$ to obtain collective QCD sum rules. The corresponding scalar invariant components are expressed as follows:

$$\begin{aligned} \Pi_{\bar{D}_0 D^* AA}(p'^2, p^2, q^2) &= \frac{\lambda_{\bar{D}_0 D^* AA}}{(m_Y^2 - p'^2)(m_{D_0}^2 - p^2)(m_{D^*}^2 - q^2)} \\ &+ \frac{C_{\bar{D}_0 D^* AA}}{(m_{D_0}^2 - p^2)(m_{D^*}^2 - q^2)} \\ &+ \frac{\bar{\lambda}_{\bar{D}_0 D^* AA}}{(m_X^2 - p'^2)(m_{D_0}^2 - p^2)(m_{D^*}^2 - q^2)} + \dots, \end{aligned} \quad (40)$$

$$\begin{aligned} \Pi_{\bar{D}_1 DAA}(p'^2, p^2, q^2) &= \frac{\lambda_{\bar{D}_1 DAA}}{(m_Y^2 - p'^2)(m_{D_1}^2 - p^2)(m_D^2 - q^2)} \\ &+ \frac{C_{\bar{D}_1 DAA}}{(m_{D_1}^2 - p^2)(m_D^2 - q^2)} \\ &+ \frac{\bar{\lambda}_{\bar{D}_1 DAA}}{(m_X^2 - p'^2)(m_{D_1}^2 - p^2)(m_D^2 - q^2)} + \dots, \end{aligned} \quad (41)$$

$$\begin{aligned} \Pi_{\chi_{c0} \omega AA}(p'^2, p^2, q^2) &= \frac{\lambda_{\chi_{c0} \omega AA}}{(m_Y^2 - p'^2)(m_{\chi_{c0}}^2 - p^2)(m_\omega^2 - q^2)} \\ &+ \frac{C_{\chi_{c0} \omega AA}}{(m_{\chi_{c0}}^2 - p^2)(m_\omega^2 - q^2)} \\ &+ \frac{\bar{\lambda}_{\chi_{c0} \omega AA}}{(m_X^2 - p'^2)(m_{\chi_{c0}}^2 - p^2)(m_\omega^2 - q^2)} + \dots, \end{aligned} \quad (42)$$

$$\begin{aligned} \Pi_{J/\psi f_0(500) AA}(p'^2, p^2, q^2) &= \frac{\lambda_{J/\psi f_0(500) AA}}{(m_Y^2 - p'^2)(m_{J/\psi}^2 - p^2)(m_{f_0(500)}^2 - q^2)} \\ &+ \frac{C_{J/\psi f_0(500) AA}}{(m_{J/\psi}^2 - p^2)(m_{f_0(500)}^2 - q^2)} \\ &+ \frac{\bar{\lambda}_{J/\psi f_0(500) AA}}{(m_X^2 - p'^2)(m_{J/\psi}^2 - p^2)(m_{f_0(500)}^2 - q^2)} + \dots, \end{aligned} \quad (43)$$

$$\begin{aligned} \Pi_{\eta_c \phi S\tilde{V}}(p'^2, p^2, q^2) &= \frac{\lambda_{\eta_c \phi S\tilde{V}}}{(m_Y^2 - p'^2)(m_{\eta_c}^2 - p^2)(m_\phi^2 - q^2)} \\ &+ \frac{C_{\eta_c \phi S\tilde{V}}}{(m_{\eta_c}^2 - p^2)(m_\phi^2 - q^2)} \\ &+ \frac{\bar{\lambda}_{\eta_c \phi S\tilde{V}}}{(m_X^2 - p'^2)(m_{\eta_c}^2 - p^2)(m_\phi^2 - q^2)} + \dots. \end{aligned} \quad (44)$$

In the above equations, we introduce the following notation to simplify the formulas.

$$\begin{aligned}
\lambda_{\bar{D}DPA} &= \frac{f_D^2 m_D^4}{m_c^2} \lambda_{PA} G_{\bar{D}DPA}, \\
\lambda_{\bar{D}^*DPA} &= \frac{f_{D^*} m_{D^*} f_D m_D^2}{m_c} \lambda_{PA} G_{\bar{D}^*DPA}, \\
\lambda_{\bar{D}^*D^*PA} &= f_{D^*}^2 m_{D^*}^2 \lambda_{PA} G_{\bar{D}^*D^*PA}, \\
\lambda_{\bar{D}_0D^*PA} &= f_{D_0} m_{D_0} f_{D^*} m_{D^*} \lambda_{PA} G_{\bar{D}_0D^*PA}, \\
\lambda_{\bar{D}_1DPA} &= \frac{f_{D_1} m_{D_1} f_D m_D^2}{m_c} \lambda_{PA} G_{\bar{D}_1DPA}, \quad (45)
\end{aligned}$$

$$\begin{aligned}
\lambda_{\eta_c \omega PA} &= \frac{f_{\eta_c} m_{\eta_c}^2 f_{\omega} m_{\omega}}{2m_c} \lambda_{PA} G_{\eta_c \omega PA}, \\
\lambda_{J/\psi \omega PA} &= f_{J/\psi} m_{J/\psi} f_{\omega} m_{\omega} \lambda_{PA} G_{J/\psi \omega PA} \left(1 + \frac{m_{\omega}^2}{m_Y^2} - \frac{m_{J/\psi}^2}{m_Y^2} \right), \\
\lambda_{\chi_{c0} \omega PA} &= f_{\chi_{c0}} m_{\chi_{c0}} f_{\omega} m_{\omega} \lambda_{PA} G_{\chi_{c0} \omega PA}, \\
\lambda_{\chi_{c1} \omega PA} &= f_{\chi_{c1}} m_{\chi_{c1}} f_{\omega} m_{\omega} \lambda_{PA} G_{\chi_{c1} \omega PA}, \\
\lambda_{J/\psi f_0(500) PA} &= f_{J/\psi} m_{J/\psi} f_{f_0(500)} m_{f_0(500)} \lambda_{PA} G_{J/\psi f_0(500) PA}. \quad (46)
\end{aligned}$$

With replacements similar to those for the scalar invariant components, we can obtain the corresponding notation for the currents $J_{\mu\nu}^{AA}$, J_{μ}^{AV} , and $J_{\mu\nu}^{SV}$, except for

$$\begin{aligned}
\lambda_{J/\psi \omega AA} &= f_{J/\psi} m_{J/\psi} f_{\omega} m_{\omega} \lambda_{AA} G_{J/\psi \omega AA}, \\
\lambda_{\eta_c \phi S \tilde{V}} &= \frac{f_{\eta_c} m_{\eta_c}^2 f_{\phi} m_{\phi}^3}{2m_c} \lambda_{S \tilde{V}} G_{\eta_c \phi S \tilde{V}}, \\
\lambda_{J/\psi \phi S \tilde{V}} &= f_{J/\psi} m_{J/\psi} f_{\phi} m_{\phi} \lambda_{S \tilde{V}} G_{J/\psi \phi S \tilde{V}}, \\
\lambda_{\chi_{c1} \phi S \tilde{V}} &= f_{\chi_{c1}} m_{\chi_{c1}}^3 f_{\phi} m_{\phi} \lambda_{S \tilde{V}} G_{\chi_{c1} \phi S \tilde{V}}, \quad (47)
\end{aligned}$$

and the notation for the axial-vector components is

$$\begin{aligned}
\bar{\lambda}_{\bar{D}_0 D^* AA} &= f_{D_0} m_{D_0}^3 f_{D^*} m_{D^*} \bar{\lambda}_{AA} \bar{G}_{\bar{D}_0 D^* AA}, \\
\bar{\lambda}_{\bar{D}_1 D AA} &= -\frac{f_{D_1} m_{D_1} f_D m_D^4}{m_c} \bar{\lambda}_{AA} \bar{G}_{\bar{D}_1 D AA}, \\
\bar{\lambda}_{\chi_{c0} \omega AA} &= f_{\chi_{c0}} m_{\chi_{c0}}^3 f_{\omega} m_{\omega} \bar{\lambda}_{AA} \bar{G}_{\chi_{c0} \omega AA}, \\
\bar{\lambda}_{J/\psi f_0(500) AA} &= -f_{J/\psi} m_{J/\psi} f_{f_0(500)} m_{f_0(500)}^3 \bar{\lambda}_{AA} \bar{G}_{J/\psi f_0(500) AA}, \\
\bar{\lambda}_{\eta_c \phi S \tilde{A}} &= \frac{f_{\eta_c} m_{\eta_c}^2 f_{\phi} m_{\phi}}{2m_c} \bar{\lambda}_{S \tilde{A}} \bar{G}_{\eta_c \phi S \tilde{A}}. \quad (48)
\end{aligned}$$

Applying the triple dispersion relation, we obtain the following equation:

$$\begin{aligned}
&\Pi_H(p'^2, p^2, q^2) \\
&= \int_{\Delta_s^2}^{\infty} ds' \int_{\Delta_s^2}^{\infty} ds \int_{\Delta_u^2}^{\infty} du \frac{\rho_H(s', s, u)}{(s' - p'^2)(s - p^2)(u - q^2)}, \quad (49)
\end{aligned}$$

where Δ_s^2 , Δ_s^2 , and Δ_u^2 are thresholds, and the subscript H denotes the hadronic side.

On the QCD side, we carry out the operator product expansion up to dimension-5 vacuum condensates and apply double dispersion relations to obtain,

$$\Pi_{\text{QCD}}(p'^2, p^2, q^2) = \int_{\Delta_s^2}^{\infty} ds \int_{\Delta_u^2}^{\infty} du \frac{\rho_{\text{QCD}}(p'^2, s, u)}{(s - p^2)(u - q^2)}, \quad (50)$$

as

$$\lim_{\epsilon \rightarrow 0} \frac{\text{Im} \Pi_{\text{QCD}}(s' + i\epsilon, p^2, q^2)}{\pi} = 0. \quad (51)$$

The triple dispersion relation in Eq. (49) on the hadron side cannot be matched with the double dispersion relation in Eq. (50) on the QCD side. Therefore, we should first perform the integration over ds' and then match the hadron side with the QCD side below the continuum thresholds to obtain rigorous quark-hadron duality [51, 54],

$$\begin{aligned}
&\int_{\Delta_s^2}^{s_0} ds \int_{\Delta_u^2}^{u_0} du \frac{\rho_{\text{QCD}}(p'^2, s, u)}{(s - p^2)(u - q^2)} \\
&= \int_{\Delta_s^2}^{s_0} ds \int_{\Delta_u^2}^{u_0} du \left[\int_{\Delta_s^2}^{\infty} ds' \frac{\rho_H(s', s, u)}{(s' - p'^2)(s - p^2)(u - q^2)} \right], \quad (52)
\end{aligned}$$

where s_0 and u_0 are the continuum thresholds. We introduce a set of free parameters, C , to parameterize the contributions of transitions between the higher resonances (continuum states) in the s' channel and the ground-state conventional meson pairs. We take the parameter $C_{\bar{D}DPA}$ in the $Y_{PA} \rightarrow \bar{D}D$ channel as an example.

$$C_{\bar{D}DPA} = \int_{s'_0}^{\infty} ds' \frac{\rho_H(s', m_{\bar{D}}^2, m_D^2)}{(s' - m_Y^2)(p^2 - m_{\bar{D}}^2)(q^2 - m_D^2)}, \quad (53)$$

where s'_0 is the continuum threshold parameter for the ground state, and $\rho_H(s', m_{\bar{D}}^2, m_D^2)$ is the formal hadronic spectral density for transitions between the higher resonances (continuum states) in the s' channel and the ground-state meson pairs $D\bar{D}$. It is obvious that, in the s and u channels, the hadron-side and QCD-side spectral densities have a one-to-one correspondence below the continuum thresholds s_0 and u_0 , respectively, whereas in the s' channel, there is no corresponding contribution on the QCD side. Experimentally, the spectroscopy of the hidden-charm tetraquark states has not yet been established. Although some exotic states are excellent candidates for tetraquark states, no definite conclusion can be drawn. We introduce a free parameter $C_{\bar{D}DPA}$ to parameterize the contributions involving the higher resonances (continuum states) in the s' channel, which results in

model dependence. At present, we have no choice but to accept this model dependence. For more detailed discussions, the reader is referred to Sec. 7 in Ref. [42].

Then, we set $p'^2 = p^2$ in the correlation functions

$$\frac{\lambda_{DDPA}}{m_Y^2 - m_D^2} \left[\exp\left(-\frac{m_D^2}{T^2}\right) - \exp\left(-\frac{m_Y^2}{T^2}\right) \right] \exp\left(-\frac{m_D^2}{T^2}\right) + C_{DDPA} \exp\left(-\frac{m_D^2}{T^2} - \frac{m_D^2}{T^2}\right) = \Pi_{DDPA}^{QCD}(T^2), \dots \quad (54)$$

In the following, we illustrate the QCD side of the QCD sum rules for the $Y_{PA} \rightarrow \bar{D}D$ channel as an example.

$$\begin{aligned} \Pi_{DDPA}^{QCD}(T^2) = & \frac{3m_c}{128\pi^4} \int_{m_c^2}^{s_D^0} ds \int_{m_c^2}^{s_D^0} du \left(1 - \frac{m_c^2}{s}\right)^2 \left(1 - \frac{m_c^2}{u}\right)^2 u \exp\left(-\frac{s+u}{T^2}\right) - \frac{\langle \bar{q}q \rangle}{16\pi^2} \int_{m_c^2}^{s_D^0} ds \left(1 - \frac{m_c^2}{s}\right)^2 \\ & \times (s + m_c^2) \exp\left(-\frac{s+m_c^2}{T^2}\right) - \frac{\langle \bar{q}g_s\sigma Gq \rangle}{384\pi^2} \int_{m_c^2}^{s_D^0} du \left(1 - \frac{m_c^2}{u}\right) \left(3 + \frac{5m_c^2}{u}\right) \exp\left(-\frac{u+m_c^2}{T^2}\right) + \frac{m_c^2 \langle \bar{q}g_s\sigma Gq \rangle}{64\pi^2 T^4} \\ & \times \int_{m_c^2}^{s_D^0} du \left(1 - \frac{m_c^2}{u}\right)^2 u \exp\left(-\frac{u+m_c^2}{T^2}\right) - \frac{m_c^2 \langle \bar{q}g_s\sigma Gq \rangle}{64\pi^2 T^2} \left(2 - \frac{m_c^2}{T^2}\right) \int_{m_c^2}^{s_D^0} ds \left(1 - \frac{m_c^2}{s}\right)^2 \exp\left(-\frac{s+m_c^2}{T^2}\right). \end{aligned} \quad (55)$$

Other QCD spectral densities are shown explicitly in Appendix C, while the expressions on the hadron side of the QCD sum rules have the same form.

In the calculations, we neglect the gluon condensates due to their tiny contributions [51, 54]. In addition, end-point divergences appear in some channels at the thresholds $s = m_c^2$, $s = 4m_c^2$ and $u = m_c^2$ due to the factors $s - m_c^2$, $s - 4m_c^2$ and $u - m_c^2$ in the denominators. The replacements $s - m_c^2 \rightarrow s - m_c^2 + \Delta^2$, $s - 4m_c^2 \rightarrow s - 4m_c^2 + \Delta^2$ and $u - m_c^2 \rightarrow u - m_c^2 + \Delta^2$ with $\Delta^2 = m_c^2$ are adopted to eliminate these divergences, as in our previous works [55–56].

III. NUMERICAL RESULTS AND DISCUSSIONS

The input parameters used in this work are listed in Table 3. We set $m_u = m_d = 0$ and take the $\overline{\text{MS}}$ masses $m_s(2\text{ GeV}) = (0.095 \pm 0.005)\text{ GeV}$ and $m_c(m_c) = (1.275 \pm 0.025)\text{ GeV}$ from the Particle Data Group [1]. We also take into account the energy-scale dependence of the vacuum condensates and the quark masses m_s and m_c using the renormalization group equations.

$$\langle \bar{q}q \rangle(\mu) = \langle \bar{q}q \rangle(1\text{ GeV}) \left[\frac{\alpha_s(1\text{ GeV})}{\alpha_s(\mu)} \right]^{\frac{12}{33-2n_f}},$$

$$\langle \bar{s}s \rangle(\mu) = \langle \bar{s}s \rangle(1\text{ GeV}) \left[\frac{\alpha_s(1\text{ GeV})}{\alpha_s(\mu)} \right]^{\frac{12}{33-2n_f}},$$

$$\langle \bar{q}g_s\sigma Gq \rangle(\mu) = \langle \bar{q}g_s\sigma Gq \rangle(1\text{ GeV}) \left[\frac{\alpha_s(1\text{ GeV})}{\alpha_s(\mu)} \right]^{\frac{2}{33-2n_f}},$$

$$\langle \bar{s}g_s\sigma Gs \rangle(\mu) = \langle \bar{s}g_s\sigma Gs \rangle(1\text{ GeV}) \left[\frac{\alpha_s(1\text{ GeV})}{\alpha_s(\mu)} \right]^{\frac{2}{33-2n_f}},$$

$\Pi(p'^2, p^2, q^2)$, perform the double Borel transformation with respect to the variables $P^2 = -p^2$ and $Q^2 = -q^2$, respectively, and set the Borel parameters $T_1^2 = T_2^2 = T^2$ to obtain the QCD sum rules,

$$\begin{aligned} m_c(\mu) &= m_c(m_c) \left[\frac{\alpha_s(\mu)}{\alpha_s(m_c)} \right]^{\frac{12}{33-2n_f}}, \\ m_s(\mu) &= m_s(2\text{ GeV}) \left[\frac{\alpha_s(\mu)}{\alpha_s(2\text{ GeV})} \right]^{\frac{12}{33-2n_f}}, \\ \alpha_s(\mu) &= \frac{1}{b_0 t} \left[1 - \frac{b_1}{b_0^2} \frac{\log t}{t} + \frac{b_1^2 (\log^2 t - \log t - 1) + b_0 b_2}{b_0^4 t^2} \right], \end{aligned} \quad (56)$$

where $t = \log \frac{\mu^2}{\Lambda_{\text{QCD}}^2}$, $b_0 = \frac{33-2n_f}{12\pi}$, $b_1 = \frac{153-19n_f}{24\pi^2}$, $b_2 = \frac{2857 - \frac{5033}{9}n_f + \frac{325}{27}n_f^2}{128\pi^3}$, $\Lambda_{\text{QCD}} = 210\text{ MeV}$, 292 MeV and 332 MeV for the flavor numbers $n_f = 5, 4$ and 3 , respectively [1, 65]. Specifically, we choose $n_f = 4$ for the hidden-charm tetraquark states and evolve all the input parameters to the typical energy scale $\mu = 1\text{ GeV}$.

We intend to study the exotic states in our unique scheme step by step: first, we study the mass spectrum, and then we study the two-body strong decays. When studying the mass spectrum of the hidden-charm (bottom) tetraquark (molecular) states, we compute the terms $g_s^2 \langle \bar{q}q \rangle^2$ with $q = u, d$ or s [24, 66–69]. In the full light-quark propagators, there are terms $\langle \bar{q}_j \gamma_\mu q_i \rangle$ [24, 66–69], which absorb the gluons emitted from the other quark lines to form $\langle \bar{q}_j \gamma_\mu q_i g_s D_\nu G_{\alpha\beta} \rangle$ and contribute to the four-quark condensate $g_s^2 \langle \bar{q}q \rangle^2$. The four-quark condensate $g_s^2 \langle \bar{q}q \rangle^2$ originates from the terms $\langle \bar{q} \gamma_\mu q g_s D_\eta G_{\lambda\tau} \rangle$, $\langle \bar{q}_j D_\mu^\dagger D_\nu^\dagger D_\alpha^\dagger q_i \rangle$ and $\langle \bar{q}_j D_\mu D_\nu D_\alpha q_i \rangle$ rather than from the radiative $O(\alpha_s)$ corrections to the four-quark condensate

Table 3. The input parameters used in the numerical calculations are given below, with the values of the vacuum condensates taken at the energy scale $\mu = 1 \text{ GeV}$.

Parameters	Values/GeV	Parameters	Values [57]	Parameters	Values
$m_{J/\psi}$	3.0969 [1]	m_{D_0}	2.40 GeV	f_{ω}, f_{ρ}	0.215 GeV [58]
m_{η_c}	2.9839 [1]	$m_{D_{s0}}$	2.32 GeV	$f_{f_0(500)}$	0.350 GeV [59, 60]
m_{ϕ}	1.019460 [1]	m_{D_1}	2.42 GeV	$f_{f_0(980)}$	0.180 GeV [61]
$m_{\chi_{c0}}$	3.41471 [1]	$m_{D_{s1}}$	2.46 GeV	λ_{PA}	$7.19 \times 10^{-2} \text{ GeV}^5$ [26]
$m_{\chi_{c1}}$	3.51067 [1]	f_D	0.208 GeV	λ_{AA}	$6.65 \times 10^{-2} \text{ GeV}^5$ [26]
m_D	1.86484 [1]	f_{D_s}	0.240 GeV	$\lambda_{AV}^{\sim}$	$1.23 \times 10^{-1} \text{ GeV}^5$ [27]
m_{D_s}	1.96835 [1]	f_{D^*}	0.263 GeV	$\lambda_{SV}^{\sim}$	$6.22 \times 10^{-2} \text{ GeV}^5$ [27]
m_{D^*}	2.00685 [1]	$f_{D_s^*}$	0.308 GeV	s_{ω}^0, s_{ρ}^0	$(1.2 \text{ GeV})^2$ [58]
$m_{D_s^*}$	2.1066 [1]	f_{D_0}	0.373 GeV	$s_{f_0(500)}^0$	1.0 GeV^2 [59, 60]
m_{ω}	0.78266 [1]	$f_{D_{s0}}$	0.333 GeV	$s_{f_0(980)}^0$	$(1.3 \text{ GeV})^2$ [61]
$m_{f_0(500)}$	0.550 [1]	f_{D_1}	0.332 GeV	$s_{J/\psi}^0$	$(3.6 \text{ GeV})^2$ [1, 62, 63]
$m_{f_0(980)}$	0.990 [1]	$f_{D_{s1}}$	0.345 GeV	$s_{\eta_c}^0$	$(3.5 \text{ GeV})^2$ [1, 62, 63]
$M_{Y(PA)}$	4.66 [26]	s_D^0	6.2 GeV^2	$s_{\chi_{c0}}^0$	$(3.9 \text{ GeV})^2$ [1, 62, 63]
$M_{Y(AA)}$	4.69 [26]	$s_{D_s}^0$	7.3 GeV^2	$s_{\chi_{c1}}^0$	$(4.0 \text{ GeV})^2$ [1, 62, 63]
$M_{Y(\widetilde{AV})}$	4.65 [27]	$s_{D^*}^0$	6.4 GeV^2	$\langle \bar{q}q \rangle$	$-(0.24 \pm 0.01 \text{ GeV})^3$ [52, 53, 64]
$M_{Y(S\widetilde{V})}$	4.68 [27]	$s_{D_s^*}^0$	7.5 GeV^2	$\langle \bar{s}s \rangle$	$(0.8 \pm 0.1) \langle \bar{q}q \rangle$ [52, 53, 64]
$f_{J/\psi}$	0.418 [62]	$s_{D_0}^0$	8.3 GeV^2	$\langle \bar{q}g_s \sigma Gq \rangle$	$m_0^2 \langle \bar{q}q \rangle$ [52, 53, 64]
f_{η_c}	0.387 [62]	$s_{D_{s0}}^0$	7.4 GeV^2	$\langle \bar{s}g_s \sigma Gs \rangle$	$m_0^2 \langle \bar{s}s \rangle$ [52, 53, 64]
$f_{\chi_{c0}}$	0.359 [63]	$s_{D_1}^0$	8.6 GeV^2	m_0^2	$(0.8 \pm 0.1) \text{ GeV}^2$ [52, 53, 64]
$f_{\chi_{c1}}$	0.338 [63]	$s_{D_{s1}}^0$	9.3 GeV^2	—	—

$\langle \bar{q}q \rangle^2$, where $D_\alpha = \partial_\alpha - ig_s G_\alpha$. The strong coupling constant $\alpha_s(\mu) = \frac{g_s^2(\mu)}{4\pi}$ appears at tree level and is energy-scale dependent. In fact, the contributions of such terms are tiny. In the present work, as in other works on the two-body strong decays of the hidden-charm tetraquark states, we neglect such terms because we only take into account vacuum condensates up to dimension 5 [42], but we consider the energy-scale dependence of the input parameters for consistency with our previous works [24, 66–69]. In Ref.[44], the vacuum condensates are taken at the energy scale $\mu = 1 \text{ GeV}$, while the c -quark mass is taken as the $\overline{\text{MS}}$ mass $m_c(m_c)$ or the approximate pole mass; this is another scheme for choosing the input parameters. In the two-point QCD sum rules for the hidden-charm tetraquark (molecular) states, the largest contributions do not come from the perturbative terms but from the vacuum condensates. Therefore, one has to calculate the radiative $O(\alpha_s)$ corrections to the perturbative terms and vacuum condensates, at least those to $\langle \bar{q}q \rangle$, at the same time if the next-to-leading contributions are required. Up to now, only the radiative $O(\alpha_s)$ corrections to the perturbative terms have been studied partially [70–72].

As we take the rigorous quark-hadron duality in the s

and u channels (see Eq. (52)), which correspond to the traditional mesons, it is reasonable to choose the typical energy scale $\mu = 1 \text{ GeV}$ as in the usual two-point QCD sum rules. If we choose a slightly larger energy scale, the integral ranges $m_c^2(\mu) - s_0/u_0$ and $4m_c^2(\mu) - s_0/u_0$ would be slightly larger; therefore, we expect a slightly larger hadronic coupling constant. As far as the light mesons $f_0(500)$, ω , $f_0(980)$ and $\phi(1020)$ are concerned, we prefer to choose $\mu = 1 \text{ GeV}$ as the universal energy scale.

In order to obtain flat platforms, the free parameters are fitted as follows:

$$\begin{aligned}
C_{\bar{D}_s D_s \widetilde{AV}} &= 0.0023 \text{ GeV}^5 \times T^2, \\
C_{\bar{D}_s^* D_s \widetilde{AV}} &= -0.0000035 \text{ GeV}^4 \times T^2, \\
C_{\bar{D}_s^* D_s^* \widetilde{AV}} &= 0.0004 \text{ GeV}^5 \times T^2, \\
C_{\bar{D}_{s0} D_s^* \widetilde{AV}} &= 0.00009 \text{ GeV}^6 \times T^2, \\
C_{\bar{D}_{s1} D_s \widetilde{AV}} &= 0.0126 \text{ GeV}^6 \times T^2, \\
C_{\eta_c \phi \widetilde{AV}} &= 0.00017 \text{ GeV}^4 \times T^2, \\
C_{J/\psi \phi \widetilde{AV}} &= 0.0, \\
C_{\chi_{c0} \phi \widetilde{AV}} &= 0.002 \text{ GeV}^6 \times T^2,
\end{aligned}$$

$$\begin{aligned} C_{\chi_{c1}\phi AV} &= 0.000012 \text{ GeV}^5 \times T^2, \\ C_{J/\psi f_0(980)AV} &= 0.0027 \text{ GeV}^6 \times T^2, \end{aligned} \quad (57)$$

$$\begin{aligned} C_{\bar{D}DPA} &= 0.00016 \text{ GeV}^5 \times T^2, \\ C_{\bar{D}^*DPA} &= -0.0000018 \text{ GeV}^4 \times T^2, \\ C_{\bar{D}^*D^*PA} &= 0.00013 \text{ GeV}^5 \times T^2, \\ C_{\bar{D}_0D^*PA} &= 0.0001 \text{ GeV}^6 \times T^2, \\ C_{\bar{D}_1DPA} &= 0.001 \text{ GeV}^6 \times T^2, \\ C_{\eta_c\omega PA} &= 0.00009 \text{ GeV}^4 \times T^2, \\ C_{J/\omega PA} &= 0.0, \\ C_{\chi_{c0}\omega PA} &= 0.00026 \text{ GeV}^6 \times T^2, \\ C_{\chi_{c1}\omega PA} &= 0.000016 \text{ GeV}^5 \times T^2, \\ C_{J/\psi f_0(500)PA} &= 0.00037 \text{ GeV}^6 \times T^2, \end{aligned} \quad (58)$$

$$\begin{aligned} C_{\bar{D}DAA} &= -0.0000013 \text{ GeV}^4 \times T^2, \\ C_{\bar{D}^*DAA} &= -0.0000012 \text{ GeV}^3 \times T^2, \\ C_{\bar{D}^*D^*AA} &= 0.0, \\ C_{\bar{D}_0D^*AA} &= 0.0003 \text{ GeV}^7, \\ \bar{\lambda}_{\bar{D}_0D^*AA} &= 0.0044 \text{ GeV}^7 \times T^2, \\ C_{\bar{D}_1DAA} &= 0.00055 \text{ GeV}^7, \\ \bar{\lambda}_{\bar{D}_1DAA} &= 0.0037 \text{ GeV}^7 \times T^2, \\ C_{\eta_c\omega AA} &= 0.0, \\ C_{J/\omega AA} &= 0.0, \\ C_{\chi_{c0}\omega AA} &= 0.00013 \text{ GeV}^7, \\ \bar{\lambda}_{\chi_{c0}\omega AA} &= 0.0034 \text{ GeV}^7 \times T^2, \\ C_{\chi_{c1}\omega AA} &= 0.043 \text{ GeV}^6, \\ C_{J/\psi f_0(500)AA} &= 0.000044 \text{ GeV}^7, \\ \bar{\lambda}_{J/\psi f_0(500)AA} &= 0.0011 \text{ GeV}^7 \times T^2, \end{aligned} \quad (59)$$

$$\begin{aligned} C_{\bar{D}_sD_s\tilde{V}} &= 0.000081 \text{ GeV}^4 \times T^2, \\ C_{\bar{D}_s^*D_s\tilde{V}} &= -0.0000013 \text{ GeV}^5 \times T^2, \\ C_{\bar{D}_s^*D_s^*\tilde{V}} &= 0.000025 \text{ GeV}^4 \times T^2, \\ C_{\bar{D}_{s0}D_s^*\tilde{V}} &= 0.00013 \text{ GeV}^5 \times T^2, \\ C_{\bar{D}_{s1}D_s\tilde{V}} &= 0.0007 \text{ GeV}^5 \times T^2, \\ C_{\eta_c\phi S\tilde{V}} &= 0.00015 \text{ GeV}^7, \\ \bar{\lambda}_{\eta_c\phi S\tilde{V}} &= 0.0033 \text{ GeV}^7 \times T^2, \end{aligned}$$

$$C_{J/\psi\phi S\tilde{V}} = 0.00002 \text{ GeV}^4 \times T^2,$$

$$C_{\chi_{c0}\phi S\tilde{V}} = 0.00032 \text{ GeV}^5 \times T^2,$$

$$C_{\chi_{c1}\phi S\tilde{V}} = 0.0004 \text{ GeV}^6 \times T^2,$$

$$C_{J/\psi f_0(980)S\tilde{V}} = 0.00022 \text{ GeV}^5 \times T^2. \quad (60)$$

We then obtain uniform flat Borel windows, $T_{\max}^2 - T_{\min}^2 = 1 \text{ GeV}^2$, which are presented explicitly in Table 4.

In Fig. 1, the curves of the hadronic coupling constants $G_{\bar{D}_{s0}D_s^*S\tilde{V}}$, $G_{\bar{D}_{s1}D_sS\tilde{V}}$, $G_{\bar{D}_sD_sS\tilde{V}}$, $G_{\bar{D}_s^*D_s^*S\tilde{V}}$, $G_{\chi_{c0}\phi S\tilde{V}}$ and $G_{J/\psi\phi S\tilde{V}}$ are plotted as functions of the Borel parameters T^2 over large intervals as examples. In the Borel windows, clear flat platforms appear, and thus the hadronic coupling constants can be extracted reliably.

The uncertainties of the hadronic coupling constants are analyzed routinely. They originate not only from the coupling constants but also from other input parameters. Taking the QCD sum rule for the channel $Y_{PA} \rightarrow \bar{D}D$ as an example, the uncertainties on the hadronic side can be written as $\lambda_{PA}\bar{f}_D f_D G_{\bar{D}DPA} = \bar{\lambda}_{PA}\bar{f}_D \bar{f}_D \bar{G}_{\bar{D}DPA} + \delta\lambda_{PA}\bar{f}_D f_D G_{\bar{D}DPA}$, $C_{\bar{D}DPA} = \bar{C}_{\bar{D}DPA} + \delta C_{\bar{D}DPA}$, $\dots$, where

$$\begin{aligned} \delta\lambda_{PA}\bar{f}_D f_D G_{\bar{D}DPA} &= \bar{\lambda}_{PA}\bar{f}_D \bar{f}_D \bar{G}_{\bar{D}DPA} \\ &\times \left(\frac{\delta f_D}{\bar{f}_D} + \frac{\delta f_D}{\bar{f}_D} + \frac{\delta\lambda_{PA}}{\bar{\lambda}_{PA}} + \frac{\delta G_{\bar{D}DPA}}{\bar{G}_{\bar{D}DPA}} \right), \end{aligned} \quad (61)$$

where the short overline denotes the central value. To avoid overestimating the uncertainties of the hadronic coupling constants, we approximately set $\delta C_{\bar{D}DPA} = 0$, $\frac{\delta f_D}{\bar{f}_D} = \frac{\delta f_D}{\bar{f}_D} = \frac{\delta\lambda_{PA}}{\bar{\lambda}_{PA}} = \frac{\delta G_{\bar{D}DPA}}{\bar{G}_{\bar{D}DPA}}$, $\dots$.

After taking into account the relevant uncertainties, the numerical values of the hadronic coupling constants can be obtained directly and are shown explicitly in Table 4. Thereafter, the partial decay widths are obtained directly using the formula,

$$\begin{aligned} \Gamma(Y \rightarrow FF') &= \frac{|T|^2 p(m_Y, m_F, m_{F'})}{24\pi m_Y^2}, \\ |T|^2 &= \Sigma |\langle F(p)F'(q)|Y(p') \rangle|^2, \end{aligned} \quad (62)$$

where $p(A, B, C) = \frac{\sqrt{[A^2 - (B+C)^2][A^2 - (B-C)^2]}}{2A}$, and F and F' represent the final states. The partial decay widths for the different channels are shown in Table 5.

Finally, we sum all the partial decay widths to obtain the total widths of these four tetraquark states.

$$\Gamma(Y_{PA}) = 391.1 \pm 21.4 \text{ MeV},$$

Table 4. The Borel parameters (windows) T^2 and the hadronic coupling constants G . Here, "Channels" denotes the subscripts of the hadronic coupling constants defined in Eqs. (18)–(19) and in Appendix A.

Channels	$T^2(\text{GeV}^2)$	G
$\bar{D}DAA$	4.7–5.7	$(2.04 \pm 0.09) \times 10^{-2} \text{ GeV}^{-1}$
$\bar{D}^*DAA$	4.2–5.2	$(1.92 \pm 0.07) \times 10^{-2} \text{ GeV}^{-2}$
$\bar{D}^*D^*AA$	---	0.0
$\bar{D}_0D^*AA$	4.0–5.0	1.99 ± 0.16
$\bar{D}_1DAA$	2.7–3.7	0.83 ± 0.08
$\eta_c\omega AA$	3.3–4.3	$(1.03 \pm 0.06) \times 10^{-2} \text{ GeV}^{-2}$
$J/\psi\omega AA$	---	0.0
$\chi_{c0}\omega AA$	4.9–5.9	1.79 ± 0.22
$\chi_{c1}\omega AA$	3.8–4.8	$(3.27 \pm 0.1) \times 10^{-1} \text{ GeV}^{-1}$
$J/\psi f_0(500)AA$	3.0–4.0	0.36 ± 0.07
$\bar{D}_sD_s\tilde{A}V$	2.2–3.2	3.35 ± 0.27
$\bar{D}_s^*D_s\tilde{A}V$	4.8–5.8	$(2.52 \pm 0.11) \times 10^{-2} \text{ GeV}^{-1}$
$\bar{D}_s^*D_s^*\tilde{A}V$	4.1–5.1	1.83 ± 0.13
$\bar{D}_{s0}D_s^*\tilde{A}V$	3.5–4.5	$0.29 \pm 0.02 \text{ GeV}^{-1}$
$\bar{D}_{s1}D_s\tilde{A}V$	1.8–2.8	$6.34 \pm 0.65 \text{ GeV}$
$\eta_c\phi\tilde{A}V$	2.3–3.3	$0.61 \pm 0.05 \text{ GeV}^{-1}$
$J/\psi\phi\tilde{A}V$	---	0.0
$\chi_{c0}\phi\tilde{A}V$	3.0–4.0	$4.02 \pm 0.47 \text{ GeV}$
$\chi_{c1}\phi\tilde{A}V$	4.7–5.7	$0.24 \pm 0.01 \text{ GeV}^{-2}$
$J/\psi f_0(980)\tilde{A}V$	1.8–2.8	$6.19 \pm 0.78 \text{ GeV}$
$\bar{D}DPA$	2.3–3.3	1.13 ± 0.08
$\bar{D}^*DPA$	4.0–5.0	$(2.45 \pm 0.10) \times 10^{-2} \text{ GeV}^{-1}$
$\bar{D}^*D^*PA$	4.1–5.1	0.34 ± 0.08
$\bar{D}_0D^*PA$	4.9–5.9	$1.25 \pm 0.06 \text{ GeV}^{-1}$
$\bar{D}_1DPA$	3.0–4.0	$3.23 \pm 0.31 \text{ GeV}$
$\eta_c\omega PA$	2.2–3.2	$0.69 \pm 0.05 \text{ GeV}$
$J/\psi\omega PA$	---	0.0
$\chi_{c0}\omega PA$	2.6–3.6	$1.41 \pm 0.19 \text{ GeV}$
$\chi_{c1}\omega PA$	5.0–6.0	$0.38 \pm 0.02 \text{ GeV}^{-2}$
$J/\psi f_0(500)PA$	1.8–2.8	$1.05 \pm 0.18 \text{ GeV}$
$\bar{D}_sD_s\tilde{S}\tilde{V}$	2.6–3.6	$0.52 \pm 0.03 \text{ GeV}^{-1}$
$\bar{D}_s^*D_s\tilde{S}\tilde{V}$	4.6–5.6	$(1.35 \pm 0.05) \times 10^{-2} \text{ GeV}^{-2}$
$\bar{D}_s^*D_s^*\tilde{S}\tilde{V}$	3.3–4.3	$0.28 \pm 0.02 \text{ GeV}^{-1}$
$\bar{D}_{s0}D_s^*\tilde{S}\tilde{V}$	4.7–5.7	3.16 ± 0.19
$\bar{D}_{s1}D_s\tilde{S}\tilde{V}$	3.9–4.9	3.02 ± 0.21
$\eta_c\phi\tilde{S}\tilde{V}$	4.6–5.6	$0.73 \pm 0.13 \text{ GeV}^{-2}$
$J/\psi\phi\tilde{S}\tilde{V}$	3.8–4.8	$0.22 \pm 0.02 \text{ GeV}^{-1}$
$\chi_{c0}\phi\tilde{S}\tilde{V}$	4.7–5.7	2.49 ± 0.23
$\chi_{c1}\phi\tilde{S}\tilde{V}$	2.7–3.7	$0.22 \pm 0.02 \text{ GeV}^{-1}$
$J/\psi f_0(980)\tilde{S}\tilde{V}$	2.3–3.3	1.36 ± 0.17




Fig. 1. (color online) The hadronic coupling constants as functions of the Borel parameters, where A , B , C , D , E , and F denote the hadronic coupling constants $G_{\bar{D}_{s0}D_s^*S\tilde{V}}$, $G_{\bar{D}_{s1}D_s^*S\tilde{V}}$, $G_{\bar{D}_sD_s^*S\tilde{V}}$, $G_{\bar{D}_s^*D_s^*S\tilde{V}}$, $G_{\chi_{c0}\phi S\tilde{V}}$, and $G_{J/\psi\phi S\tilde{V}}$, respectively.

$$\begin{aligned}
 \Gamma(Y_{AA}) &= 27.8 \pm 2.3 \text{ MeV}, \\
 \Gamma(Y_{AV}) &= 266.7 \pm 23.9 \text{ MeV}, \\
 \Gamma(Y_{S\tilde{V}}) &= 61.5 \pm 7.3 \text{ MeV}.
 \end{aligned} \tag{63}$$

It can be clearly seen that the values of the total widths of the four states are quite different from each other. Therefore, we can distinguish them easily in high-energy experiments. The predicted width $61.5 \pm 7.3 \text{ MeV}$ of the $Y_{S\tilde{V}}$ is well compatible with the width $48 \pm 15 \pm 3 \text{ MeV}$ reported by the Belle collaboration [2]. Moreover, our prediction for the width of the $Y_{S\tilde{V}}$ is in excellent agreement with the average width $55 \pm 9 \text{ MeV}$ of the $Y(4660)$ from different experiments reported by the Particle Data Group [1], which supports the assignment of the $Y(4660)$ as the $[sc]_s[\bar{s}\bar{c}]_{\tilde{V}} - [sc]_{\tilde{V}}[\bar{s}\bar{c}]_s$ tetraquark state. The widths $155.4 \pm 24.8 \pm 0.8$ and 218.3 ± 71.9 [5] reported by the BE-

SIII collaboration are larger than those from the Belle and BaBar collaborations, which suggests that the $Y(4660)$ may have several Fock components. The predictions $\Gamma(Y_{AV}) = 266.7 \pm 23.9 \text{ MeV}$ and $\Gamma(Y_{PA}) = 391.1 \pm 21.4 \text{ MeV}$ are too large, whereas the prediction $\Gamma(Y_{AA}) = 27.8 \pm 2.3 \text{ MeV}$ is too small to be consistent with the experimental width. Thus, the present study disfavors the assignment of the $Y(4660)$ as the $[sc]_{\tilde{A}}[\bar{s}\bar{c}]_V + [sc]_V[\bar{s}\bar{c}]_{\tilde{A}}$, $[uc]_P[\bar{u}\bar{c}]_A + [dc]_P[\bar{d}\bar{c}]_A - [uc]_A[\bar{u}\bar{c}]_P - [dc]_A[\bar{d}\bar{c}]_P$, or $[uc]_A[\bar{u}\bar{c}]_A + [dc]_A[\bar{d}\bar{c}]_A$ tetraquark states. These results are useful for diagnosing vector exotic states and await future experimental examination.

IV. CONCLUSION

In this work, we consider four four-quark currents to explore the two-body strong decays of the

Table 5. Partial decay widths of the vector tetraquark states Y_{AA} , Y_{AV} , Y_{PA} , and Y_{SV} .

Channels	Γ/MeV
$Y_{AA} \rightarrow \bar{D}^0 D^0, \bar{D}^- D^+$	0.003 ± 0.0
$Y_{AA} \rightarrow \frac{\bar{D}^{*0} D^0 + \bar{D}^0 D^{*0}}{\sqrt{2}}, \frac{\bar{D}^{*-} D^+ + \bar{D}^- D^{*+}}{\sqrt{2}}$	0.023 ± 0.002
$Y_{AA} \rightarrow \bar{D}^{*0} D^{*0}, \bar{D}^{*-} D^{*+}$	0.0
$Y_{AA} \rightarrow \frac{\bar{D}_0^0 D^{*0} + \bar{D}^{*0} D_0^0}{\sqrt{2}}, \frac{\bar{D}_0^- D^{*+} + \bar{D}^{*-} D_0^+}{\sqrt{2}}$	6.03 ± 0.97
$Y_{AA} \rightarrow \frac{\bar{D}_1^0 D^0 - \bar{D}^0 D_1^0}{\sqrt{2}}, \frac{\bar{D}_1^- D^+ - \bar{D}^- D_1^+}{\sqrt{2}}$	1.23 ± 0.25
$Y_{AA} \rightarrow \eta_c \omega$	0.005 ± 0.0
$Y_{AA} \rightarrow J/\psi \omega$	0.0
$Y_{AA} \rightarrow \chi_{c0} \omega$	7.07 ± 1.74
$Y_{AA} \rightarrow \chi_{c1} \omega$	5.81 ± 0.36
$Y_{AA} \rightarrow J/\psi f_0(500)$	0.31 ± 0.12
$Y_{AV} \rightarrow \bar{D}_s D_s$	52.04 ± 8.39
$Y_{AV} \rightarrow \frac{\bar{D}_s^* D_s + \bar{D}_s D_s^*}{\sqrt{2}}$	0.023 ± 0.002
$Y_{AV} \rightarrow \bar{D}_s^* D_s^*$	30.35 ± 4.31
$Y_{AV} \rightarrow \frac{\bar{D}_{s0} D_s^* + \bar{D}_s^* D_{s0}}{\sqrt{2}}$	3.96 ± 0.57
$Y_{AV} \rightarrow \frac{\bar{D}_{s1} D_s - \bar{D}_s D_{s1}}{\sqrt{2}}$	53.52 ± 10.97
$Y_{AV} \rightarrow \eta_c \phi$	12.24 ± 1.92
$Y_{AV} \rightarrow J/\psi \phi$	0.0
$Y_{AV} \rightarrow \chi_{c0} \phi$	19.91 ± 4.65
$Y_{AV} \rightarrow \chi_{c1} \phi$	22.56 ± 2.23
$Y_{AV} \rightarrow J/\psi f_0(980)$	72.08 ± 18.16
$Y_{PA} \rightarrow \bar{D}^0 D^0, \bar{D}^- D^+$	8.50 ± 1.20
$Y_{PA} \rightarrow \frac{\bar{D}^{*0} D^0 + \bar{D}^0 D^{*0}}{\sqrt{2}}, \frac{\bar{D}^{*-} D^+ + \bar{D}^- D^{*+}}{\sqrt{2}}$	0.035 ± 0.003
$Y_{PA} \rightarrow \bar{D}^{*0} D^{*0}, \bar{D}^{*-} D^{*+}$	2.26 ± 1.00
$Y_{PA} \rightarrow \frac{\bar{D}_0^0 D^{*0} + \bar{D}^{*0} D_0^0}{\sqrt{2}}, \frac{\bar{D}_0^- D^{*+} + \bar{D}^{*-} D_0^+}{\sqrt{2}}$	80.49 ± 7.73
$Y_{PA} \rightarrow \frac{\bar{D}_1^0 D^0 - \bar{D}^0 D_1^0}{\sqrt{2}}, \frac{\bar{D}_1^- D^+ - \bar{D}^- D_1^+}{\sqrt{2}}$	18.20 ± 3.49
$Y_{PA} \rightarrow \eta_c \omega$	22.52 ± 3.51
$Y_{PA} \rightarrow J/\psi \omega$	0.0
$Y_{PA} \rightarrow \chi_{c0} \omega$	4.20 ± 1.13
$Y_{PA} \rightarrow \chi_{c1} \omega$	142.85 ± 17.20
$Y_{PA} \rightarrow J/\psi f_0(500)$	2.58 ± 0.88
$Y_{SV} \rightarrow \bar{D}_s D_s$	1.34 ± 0.14
$Y_{SV} \rightarrow \frac{\bar{D}_s^* D_s + \bar{D}_s D_s^*}{\sqrt{2}}$	0.007 ± 0.001
$Y_{SV} \rightarrow \bar{D}_s^* D_s^*$	0.80 ± 0.09
$Y_{SV} \rightarrow \frac{\bar{D}_{s0} D_s^* + \bar{D}_s^* D_{s0}}{\sqrt{2}}$	14.19 ± 1.71
$Y_{SV} \rightarrow \frac{\bar{D}_{s1} D_s - \bar{D}_s D_{s1}}{\sqrt{2}}$	12.85 ± 1.79
$Y_{SV} \rightarrow \eta_c \phi$	18.71 ± 6.63
$Y_{SV} \rightarrow J/\psi \phi$	0.60 ± 0.08
$Y_{SV} \rightarrow \chi_{c0} \phi$	8.20 ± 1.52
$Y_{SV} \rightarrow \chi_{c1} \phi$	1.24 ± 0.24
$Y_{SV} \rightarrow J/\psi f_0(980)$	3.54 ± 0.89

$[sc]_A[\bar{s}\bar{c}]_V + [sc]_V[\bar{s}\bar{c}]_A, [sc]_S[\bar{s}\bar{c}]_V - [sc]_V[\bar{s}\bar{c}]_S, [uc]_P[\bar{u}\bar{c}]_A + [dc]_P[\bar{d}\bar{c}]_A - [uc]_A[\bar{u}\bar{c}]_P - [dc]_A[\bar{d}\bar{c}]_P$ and $[uc]_A[\bar{u}\bar{c}]_A + [dc]_A[\bar{d}\bar{c}]_A$ tetraquark states around 4.66 GeV, all with the quantum numbers $J^{PC} = 1^{--}$, within the framework of QCD sum rules. We perform the operator product expansion including vacuum condensates up to dimension 5 and match the QCD side with the hadronic side based on rigorous quark-hadron duality. The resulting total widths of these states are quite different from one another. The predicted width of 61.5 ± 7.3 MeV for the $Y_{S\bar{V}}$ is in excellent agreement with the experimental values for the $Y(4660)$, favoring the $[sc]_S[\bar{s}\bar{c}]_V - [sc]_V[\bar{s}\bar{c}]_S$ -type tetraquark interpretation of the $Y(4660)$. The predictions for the other tetraquark states serve as a guide for future experiments.

APPENDIX

A. The hadronic coupling constants

In this section, we present the definitions of the other hadronic coupling constants.

$$\begin{aligned} \langle \bar{D}^*(p) D^*(q) | Y_{PA}(p') \rangle &= i\xi^* \cdot \xi^*(p-q) \cdot \varepsilon G_{\bar{D}^* D^* PA}, \\ \langle \bar{D}_s^*(p) D_s^*(q) | Y_{AV}(p') \rangle &= -\xi^* \cdot \xi^*(p-q) \cdot \varepsilon G_{\bar{D}_s^* D_s^* AV}, \\ \langle \bar{D}^*(p) D^*(q) | Y_{AA}(p') \rangle &= i\xi^* \cdot \xi^*(p-q) \cdot \varepsilon G_{\bar{D}^* D^* AA}, \\ \langle \bar{D}_s^*(p) D_s^*(q) | Y_{S\bar{V}}(p') \rangle &= -i\xi^* \cdot \xi^*(p-q) \cdot \varepsilon G_{\bar{D}_s^* D_s^* S\bar{V}}, \end{aligned} \quad (A1)$$

$$\begin{aligned} \langle \bar{D}_0(p) D^*(q) | Y_{PA}(p') \rangle &= -i\xi^* \cdot \varepsilon p \cdot q G_{\bar{D}_0 D^* PA}, \\ \langle \bar{D}_{s0}(p) D_s^*(q) | Y_{AV}(p') \rangle &= \xi^* \cdot \varepsilon G_{\bar{D}_{s0} D_s^* AV}, \\ \langle \bar{D}_0(p) D^*(q) | Y_{AA}(p') \rangle &= i\xi^* \cdot \varepsilon p \cdot q G_{\bar{D}_0 D^* AA}, \\ \langle \bar{D}_0(p) D^*(q) | X_{AA}(p') \rangle &= i\varepsilon^{\lambda\tau\rho\sigma} q_\lambda \xi_\tau^* p'_\rho \varepsilon_\sigma \bar{G}_{\bar{D}_0 D^* AA}, \\ \langle \bar{D}_{s0}(p) D_s^*(q) | Y_{S\bar{V}}(p') \rangle &= -i\xi^* \cdot \varepsilon G_{\bar{D}_{s0} D_s^* S\bar{V}}, \end{aligned} \quad (A2)$$

$$\begin{aligned} \langle \bar{D}_1(p) D(q) | Y_{PA}(p') \rangle &= -\xi^* \cdot \varepsilon G_{\bar{D}_1 D PA}, \\ \langle \bar{D}_{s1}(p) D_s(q) | Y_{AV}(p') \rangle &= i\xi^* \cdot \varepsilon G_{\bar{D}_{s1} D_s AV}, \\ \langle \bar{D}_1(p) D(q) | Y_{AA}(p') \rangle &= -i\xi^* \cdot \varepsilon G_{\bar{D}_1 D AA}, \\ \langle \bar{D}_1(p) D(q) | X_{AA}(p') \rangle &= i\varepsilon^{\lambda\tau\rho\sigma} p_\lambda \xi_\tau^* p'_\rho \varepsilon_\sigma \bar{G}_{\bar{D}_1 D AA}, \\ \langle \bar{D}_{s1}(p) D_s(q) | Y_{S\bar{V}}(p') \rangle &= -\xi^* \cdot \varepsilon G_{\bar{D}_{s1} D_s S\bar{V}}, \end{aligned} \quad (A3)$$

$$\begin{aligned} \langle \eta_c(p) \omega(q) | Y_{PA}(p') \rangle &= i\varepsilon^{\lambda\tau\rho\sigma} q_\lambda \xi_\tau^* p'_\rho \varepsilon_\sigma G_{\eta_c \omega PA}, \\ \langle \eta_c(p) \phi(q) | Y_{AV}(p') \rangle &= i\varepsilon^{\lambda\tau\rho\sigma} q_\lambda \xi_\tau^* p'_\rho \varepsilon_\sigma G_{\eta_c \phi AV}, \\ \langle \eta_c(p) \omega(q) | Y_{AA}(p') \rangle &= -i\varepsilon^{\lambda\tau\rho\sigma} q_\lambda \xi_\tau^* p'_\rho \varepsilon_\sigma G_{\eta_c \omega AA}, \\ \langle \eta_c(p) \phi(q) | Y_{S\bar{V}}(p') \rangle &= i\varepsilon^{\lambda\tau\rho\sigma} q_\lambda \xi_\tau^* p'_\rho \varepsilon_\sigma G_{\eta_c \phi S\bar{V}}, \end{aligned}$$

$$\langle \eta_c(p) \phi(q) | X_{SA}(p') \rangle = i\xi^* \cdot \varepsilon \bar{G}_{\eta_c \phi SA}, \quad (A4)$$

$$\begin{aligned} \langle J/\psi(p) \omega(q) | Y_{PA}(p') \rangle &= \xi^* \cdot \xi^*(p-q) \cdot \varepsilon G_{J/\psi \omega PA}, \\ \langle J/\psi(p) \phi(q) | Y_{AV}(p') \rangle &= \xi^* \cdot \xi^*(p-q) \cdot \varepsilon G_{J/\psi \phi AV}, \\ \langle J/\psi(p) \omega(q) | Y_{AA}(p') \rangle &= i\xi^* \cdot \xi^*(p-q) \cdot \varepsilon G_{J/\psi \omega AA}, \\ \langle J/\psi(p) \phi(q) | Y_{S\bar{V}}(p') \rangle &= -i\xi^* \cdot \xi^*(p-q) \cdot \varepsilon G_{J/\psi \phi S\bar{V}}, \end{aligned} \quad (A5)$$

$$\begin{aligned} \langle \chi_{c0}(p) \omega(q) | Y_{PA}(p') \rangle &= i\xi^* \cdot \varepsilon G_{\chi_{c0} \omega PA}, \\ \langle \chi_{c0}(p) \phi(q) | Y_{AV}(p') \rangle &= -\xi^* \cdot \varepsilon G_{\chi_{c0} \phi AV}, \\ \langle \chi_{c0}(p) \omega(q) | Y_{AA}(p') \rangle &= i\xi^* \cdot \varepsilon G_{\chi_{c0} \omega AA}, \\ \langle \chi_{c0}(p) \omega(q) | X_{AA}(p') \rangle &= i\varepsilon^{\lambda\tau\rho\sigma} q_\lambda \xi_\tau^* p'_\rho \varepsilon_\sigma \bar{G}_{\chi_{c0} \omega AA}, \\ \langle \chi_{c0}(p) \phi(q) | Y_{S\bar{V}}(p') \rangle &= -i\xi^* \cdot \varepsilon G_{\chi_{c0} \phi S\bar{V}}, \end{aligned} \quad (A6)$$

$$\begin{aligned} \langle \chi_{c1}(p) \omega(q) | Y_{PA}(p') \rangle &= \varepsilon^{\rho\sigma\lambda\tau} p_\rho \xi_\sigma^* \xi_\lambda^* \varepsilon_\tau p \cdot q G_{\chi_{c1} \omega PA}, \\ \langle \chi_{c1}(p) \phi(q) | Y_{AV}(p') \rangle &= -i\varepsilon^{\rho\sigma\lambda\tau} p_\rho \xi_\sigma^* \xi_\lambda^* \varepsilon_\tau p \cdot q G_{\chi_{c1} \phi AV}, \\ \langle \chi_{c1}(p) \omega(q) | Y_{AA}(p') \rangle &= -\varepsilon^{\rho\sigma\lambda\tau} p_\rho \xi_\sigma^* \xi_\lambda^* \varepsilon_\tau p \cdot q G_{\chi_{c1} \omega AA}, \\ \langle \chi_{c1}(p) \phi(q) | Y_{S\bar{V}}(p') \rangle &= \varepsilon^{\rho\sigma\lambda\tau} p_\rho \xi_\sigma^* \xi_\lambda^* \varepsilon_\tau G_{\chi_{c1} \phi S\bar{V}}, \end{aligned} \quad (A7)$$

$$\begin{aligned} \langle J/\psi(p) f_0(500)(q) | Y_{PA}(p') \rangle &= -i\xi^* \cdot \varepsilon G_{J/\psi f_0(500) PA}, \\ \langle J/\psi(p) f_0(980)(q) | Y_{AV}(p') \rangle &= \xi^* \cdot \varepsilon G_{J/\psi f_0(980) AV}, \\ \langle J/\psi(p) f_0(500)(q) | Y_{AA}(p') \rangle &= i\xi^* \cdot \varepsilon G_{J/\psi f_0(500) AA}, \\ \langle J/\psi(p) f_0(500)(q) | X_{AA}(p') \rangle &= i\varepsilon^{\rho\sigma\lambda\tau} p_\rho \xi_\sigma^* p'_\lambda \varepsilon_\tau \bar{G}_{J/\psi f_0(500) AA}, \\ \langle J/\psi(p) f_0(980)(q) | Y_{S\bar{V}}(p') \rangle &= -i\xi^* \cdot \varepsilon G_{J/\psi f_0(980) S\bar{V}}. \end{aligned} \quad (A8)$$

B. The correlation functions

In this section, we present the phenomenological-side correlation functions for the currents $J_{\mu\nu}^{AA}$, J_{μ}^{AV} , and $J_{\mu\nu}^{SV}$.

$$\Pi_{\mu\nu}^{\bar{D}^{AA}}(p, q) = \Pi_{\bar{D}^{AA}}(p'^2, p^2, q^2) [2(p_\mu q_\nu - p_\nu q_\mu)] + \cdots, \quad (B1)$$

$$\begin{aligned} \Pi_{\alpha\mu\nu}^{\bar{D}^* D^{AA}}(p, q) &= \Pi_{\bar{D}^* D^{AA}}(p'^2, p^2, q^2) \left(\varepsilon_{\alpha\mu\lambda\tau} p^\lambda q^\tau q_\nu \right. \\ &\quad \left. - \varepsilon_{\alpha\nu\lambda\tau} p^\lambda q^\tau q_\mu \right) + \cdots, \end{aligned} \quad (B2)$$

$$\Pi_{\alpha\beta\mu\nu}^{\bar{D}^* D^* AA}(p, q) = \Pi_{\bar{D}^* D^* AA}(p'^2, p^2, q^2) [2g_{\alpha\beta}(p_\mu q_\nu - p_\nu q_\mu)] + \cdots, \quad (B3)$$

$$\Pi_{\alpha\mu\nu}^{\bar{D}_0 D^* AA}(p, q) = \Pi_{\bar{D}_0 D^* AA}(p'^2, p^2, q^2) (g_{\alpha\mu} q_\nu) + \cdots, \quad (\text{B4})$$

$$\Pi_{\alpha\mu\nu}^{\bar{D}_1 D^* AA}(p, q) = \Pi_{\bar{D}_1 D^* AA}(p'^2, p^2, q^2) (-g_{\alpha\mu} p_\nu) + \cdots, \quad (\text{B5})$$

$$\Pi_{\alpha\mu\nu}^{\eta_c \omega AA}(p, q) = \Pi_{\eta_c \omega AA}(p'^2, p^2, q^2) (\varepsilon_{\alpha\mu\lambda\tau} p^\lambda p_\nu q^\tau - \varepsilon_{\alpha\nu\lambda\tau} p^\lambda p_\mu q^\tau) + \cdots, \quad (\text{B6})$$

$$\Pi_{\alpha\beta\mu\nu}^{J/\psi \omega AA}(p, q) = \Pi_{J/\psi \omega AA}(p'^2, p^2, q^2) [2g_{\alpha\beta}(p_\mu q_\nu - p_\nu q_\mu)] + \cdots, \quad (\text{B7})$$

$$\Pi_{\alpha\mu\nu}^{\chi_{c0} \omega AA}(p, q) = \Pi_{\chi_{c0} \omega AA}(p'^2, p^2, q^2) (g_{\alpha\mu} q_\nu) + \cdots, \quad (\text{B8})$$

$$\Pi_{\alpha\beta\mu\nu}^{\chi_{c1} \omega AA}(p, q) = \Pi_{\chi_{c1} \omega AA}(p'^2, p^2, q^2) i (\varepsilon_{\alpha\beta\mu\tau} p^\tau q_\nu - \varepsilon_{\alpha\beta\nu\tau} p^\tau q_\mu) + \cdots, \quad (\text{B9})$$

$$\Pi_{\alpha\mu\nu}^{J/\psi f_0(500) AA}(p, q) = \Pi_{J/\psi f_0(500) AA}(p'^2, p^2, q^2) (g_{\alpha\mu} p_\nu) + \cdots, \quad (\text{B10})$$

$$\Pi_{\mu}^{\bar{D}_s D_s \tilde{A} V}(p, q) = \Pi_{\bar{D}_s D_s \tilde{A} V}(p'^2, p^2, q^2) i(p - q)_\mu + \cdots, \quad (\text{B11})$$

$$\Pi_{\alpha\mu}^{\bar{D}_s^* D_s \tilde{A} V}(p, q) = \Pi_{\bar{D}_s^* D_s \tilde{A} V}(p'^2, p^2, q^2) (-i\varepsilon_{\alpha\mu\lambda\tau} p^\lambda q^\tau) + \cdots, \quad (\text{B12})$$

$$\Pi_{\alpha\beta\mu}^{\bar{D}_s^* D_s \tilde{A} V}(p, q) = \Pi_{\bar{D}_s^* D_s \tilde{A} V}(p'^2, p^2, q^2) (-ig_{\alpha\beta} p_\mu) + \cdots, \quad (\text{B13})$$

$$\Pi_{\alpha\mu}^{\bar{D}_{s0} D_s \tilde{A} V}(p, q) = \Pi_{\bar{D}_{s0} D_s \tilde{A} V}(p'^2, p^2, q^2) (-ig_{\alpha\mu} p \cdot q) + \cdots, \quad (\text{B14})$$

$$\Pi_{\alpha\mu}^{\bar{D}_{s1} D_s \tilde{A} V}(p, q) = \Pi_{\bar{D}_{s1} D_s \tilde{A} V}(p'^2, p^2, q^2) (g_{\alpha\mu}) + \cdots, \quad (\text{B15})$$

$$\Pi_{\alpha\mu}^{\eta_c \phi \tilde{A} V}(p, q) = \Pi_{\eta_c \phi \tilde{A} V}(p'^2, p^2, q^2) (-i\varepsilon_{\alpha\mu\lambda\tau} p^\lambda q^\tau) + \cdots, \quad (\text{B16})$$

$$\Pi_{\alpha\beta\mu}^{J/\psi \phi \tilde{A} V}(p, q) = \Pi_{J/\psi \phi \tilde{A} V}(p'^2, p^2, q^2) (ig_{\alpha\beta} p_\mu) + \cdots, \quad (\text{B17})$$

$$\Pi_{\alpha\mu}^{\chi_{c0} \phi \tilde{A} V}(p, q) = \Pi_{\chi_{c0} \phi \tilde{A} V}(p'^2, p^2, q^2) (ig_{\alpha\mu}) + \cdots, \quad (\text{B18})$$

$$\Pi_{\alpha\beta\mu}^{\chi_{c1} \phi \tilde{A} V}(p, q) = \Pi_{\chi_{c1} \phi \tilde{A} V}(p'^2, p^2, q^2) (-\varepsilon_{\alpha\beta\mu\tau} p^\tau p \cdot q) + \cdots, \quad (\text{B19})$$

$$\Pi_{\alpha\mu}^{J/\psi f_0(980) \tilde{A} V}(p, q) = \Pi_{J/\psi f_0(980) \tilde{A} V}(p'^2, p^2, q^2) (-ig_{\alpha\mu}) + \cdots, \quad (\text{B20})$$

$$\Pi_{\mu\nu}^{\bar{D}_s D_s \tilde{S} V}(p, q) = \Pi_{\bar{D}_s D_s \tilde{S} V}(p'^2, p^2, q^2) (-2\varepsilon_{\mu\nu\lambda\tau} p^\lambda q^\tau) + \cdots, \quad (\text{B21})$$

$$\Pi_{\alpha\mu\nu}^{\bar{D}_s^* D_s \tilde{S} V}(p, q) = \Pi_{\bar{D}_s^* D_s \tilde{S} V}(p'^2, p^2, q^2) p \cdot q [g_{\mu\alpha}(p - q)_\nu - g_{\nu\alpha}(p - q)_\mu] + \cdots, \quad (\text{B22})$$

$$\Pi_{\alpha\beta\mu\nu}^{\bar{D}_s^* D_s \tilde{S} V}(p, q) = \Pi_{\bar{D}_s^* D_s \tilde{S} V}(p'^2, p^2, q^2) (-2g_{\alpha\beta} \varepsilon_{\mu\nu\lambda\tau} p^\lambda q^\tau) + \cdots, \quad (\text{B23})$$

$$\Pi_{\alpha\mu\nu}^{\bar{D}_{s0} D_s \tilde{S} V}(p, q) = \Pi_{\bar{D}_{s0} D_s \tilde{S} V}(p'^2, p^2, q^2) (-\varepsilon_{\alpha\mu\nu\tau} p^\tau) + \cdots, \quad (\text{B24})$$

$$\Pi_{\alpha\mu\nu}^{\bar{D}_{s1} D_s \tilde{S} V}(p, q) = \Pi_{\bar{D}_{s1} D_s \tilde{S} V}(p'^2, p^2, q^2) (i\varepsilon_{\alpha\mu\nu\tau} q^\tau) + \cdots, \quad (\text{B25})$$

$$\Pi_{\alpha\mu\nu}^{\eta_c \phi \tilde{S} V}(p, q) = \Pi_{\eta_c \phi \tilde{S} V}(p'^2, p^2, q^2) (g_{\mu\alpha} p_\nu - g_{\nu\alpha} p_\mu) + \cdots, \quad (\text{B26})$$

$$\Pi_{\alpha\beta\mu\nu}^{J/\psi \phi \tilde{S} V}(p, q) = \Pi_{J/\psi \phi \tilde{S} V}(p'^2, p^2, q^2) (2g_{\alpha\beta} \varepsilon_{\mu\nu\lambda\tau} p^\lambda q^\tau) + \cdots, \quad (\text{B27})$$

$$\Pi_{\alpha\mu\nu}^{\chi_{c0} \phi \tilde{S} V}(p, q) = \Pi_{\chi_{c0} \phi \tilde{S} V}(p'^2, p^2, q^2) (-\varepsilon_{\alpha\mu\nu\tau} q^\tau) + \cdots, \quad (\text{B28})$$

$$\Pi_{\alpha\beta\mu\nu}^{\chi_{c1} \omega \tilde{S} V}(p, q) = \Pi_{\chi_{c1} \omega \tilde{S} V}(p'^2, p^2, q^2) i(g_{\alpha\mu} g_{\beta\nu} - g_{\alpha\nu} g_{\beta\mu}) + \cdots, \quad (\text{B29})$$

$$\Pi_{\alpha\mu\nu}^{J/\psi f_0(980) \tilde{S} V}(p, q) = \Pi_{J/\psi f_0(980) \tilde{S} V}(p'^2, p^2, q^2) (-\varepsilon_{\alpha\mu\nu\tau} p^\tau) + \cdots. \quad (\text{B30})$$

C. The spectral densities

In this section, we present the explicit expressions for the QCD side of the QCD sum rules.

C.1. The QCD side for the current J_μ^{PA}

$$\Pi_{D^*D^{PA}}^{\text{QCD}}(T^2) = -\frac{m_c \langle \bar{q} g_s \sigma G q \rangle}{192\pi^2} \int_{m_c^2}^{s_D^0} du \left(3 - \frac{m_c^2}{u} \right) \frac{1}{u} \exp\left(-\frac{u+m_c^2}{T^2}\right) + \frac{m_c \langle \bar{q} g_s \sigma G q \rangle}{64\pi^2} \int_{m_c^2}^{s_{D^*}^0} ds \left(1 - \frac{m_c^2}{s} \right) \frac{1}{s} \exp\left(-\frac{s+m_c^2}{T^2}\right), \quad (\text{C1})$$

$$\begin{aligned} \Pi_{D^*D^{PA}}^{\text{QCD}}(T^2) &= \frac{m_c}{128\pi^4} \int_{m_c^2}^{s_{D^*}^0} ds \int_{m_c^2}^{s_{D^*}^0} du \left(1 - \frac{m_c^2}{s} \right)^2 \left(1 - \frac{m_c^2}{u} \right)^2 (2u+m_c^2) \exp\left(-\frac{s+u}{T^2}\right) + \frac{\langle \bar{q} q \rangle}{24\pi^2} \int_{m_c^2}^{s_{D^*}^0} ds \left(1 - \frac{m_c^2}{s} \right)^2 \\ &\times (s-m_c^2) \exp\left(-\frac{s+m_c^2}{T^2}\right) + \frac{\langle \bar{q} g_s \sigma G q \rangle}{576\pi^2 T^2} \left(4 + \frac{3m_c^2}{T^2} \right) \int_{m_c^2}^{s_{D^*}^0} du \left(1 - \frac{m_c^2}{u} \right)^2 (2u+m_c^2) \exp\left(-\frac{u+m_c^2}{T^2}\right) \\ &+ \frac{m_c^4 \langle \bar{q} g_s \sigma G q \rangle}{64\pi^2 T^4} \int_{m_c^2}^{s_{D^*}^0} ds \left(1 - \frac{m_c^4}{s^2} \right) \exp\left(-\frac{s+m_c^2}{T^2}\right) - \frac{m_c^2 \langle \bar{q} g_s \sigma G q \rangle}{192\pi^2} \int_{m_c^2}^{s_{D^*}^0} ds \frac{1}{s} \left(2 - \frac{4m_c^2}{s} \right) \exp\left(-\frac{s+m_c^2}{T^2}\right), \end{aligned} \quad (\text{C2})$$

$$\begin{aligned} \Pi_{D_0D^{PA}}^{\text{QCD}}(T^2) &= \frac{3m_c^2}{128\pi^4} \int_{m_c^2}^{s_{D_0}^0} ds \int_{m_c^2}^{s_{D^*}^0} du \left(1 - \frac{m_c^2}{s} \right)^2 \left(1 - \frac{m_c^4}{u^2} \right) \exp\left(-\frac{s+u}{T^2}\right) - \frac{m_c \langle \bar{q} q \rangle}{16\pi^2} \int_{m_c^2}^{s_{D_0}^0} ds \left(1 - \frac{m_c^2}{s} \right)^2 \exp\left(-\frac{s+m_c^2}{T^2}\right) \\ &+ \frac{m_c \langle \bar{q} q \rangle}{16\pi^2} \int_{m_c^2}^{s_{D^*}^0} du \left(1 - \frac{m_c^2}{u} \right)^2 \exp\left(-\frac{u+m_c^2}{T^2}\right) - \frac{m_c^3 \langle \bar{q} g_s \sigma G q \rangle}{64\pi^2 T^4} \int_{m_c^2}^{s_{D^*}^0} du \left(1 - \frac{m_c^2}{u} \right)^2 \exp\left(-\frac{u+m_c^2}{T^2}\right) \\ &+ \frac{m_c \langle \bar{q} g_s \sigma G q \rangle}{192\pi^2 T^2} \left(4 + \frac{3m_c^2}{T^2} \right) \int_{m_c^2}^{s_{D_0}^0} ds \left(1 - \frac{m_c^2}{s} \right)^2 \exp\left(-\frac{s+m_c^2}{T^2}\right) + \frac{m_c \langle \bar{q} g_s \sigma G q \rangle}{192\pi^2} \int_{m_c^2}^{s_{D^*}^0} du \left(1 + \frac{m_c^4}{u^2} \right) \frac{1}{u-m_c^2} \\ &\times \exp\left(-\frac{u+m_c^2}{T^2}\right) + \frac{m_c^3 \langle \bar{q} g_s \sigma G q \rangle}{64\pi^2} \int_{m_c^2}^{s_{D^*}^0} du \frac{1}{u^2} \exp\left(-\frac{u+m_c^2}{T^2}\right) - \frac{m_c \langle \bar{q} g_s \sigma G q \rangle}{192\pi^2} \int_{m_c^2}^{s_{D_0}^0} ds \left(3 - \frac{2m_c^2}{s} \right) \\ &\times \frac{1}{s} \exp\left(-\frac{s+m_c^2}{T^2}\right), \end{aligned} \quad (\text{C3})$$

$$\begin{aligned} \Pi_{D_1D^{PA}}^{\text{QCD}}(T^2) &= \frac{1}{128\pi^4} \int_{m_c^2}^{s_{D_1}^0} ds \int_{m_c^2}^{s_{D^*}^0} du \left(1 - \frac{m_c^2}{s} \right)^2 \left(1 - \frac{m_c^2}{u} \right)^2 u (2s+m_c^2) \exp\left(-\frac{s+u}{T^2}\right) - \frac{m_c \langle \bar{q} q \rangle}{48\pi^2} \int_{m_c^2}^{s_{D_1}^0} ds \left(1 - \frac{m_c^2}{s} \right)^2 \\ &\times (2s+m_c^2) \exp\left(-\frac{s+m_c^2}{T^2}\right) + \frac{m_c \langle \bar{q} q \rangle}{16\pi^2} \int_{m_c^2}^{s_{D^*}^0} du u \left(1 - \frac{m_c^2}{u} \right)^2 \exp\left(-\frac{u+m_c^2}{T^2}\right) \\ &- \frac{m_c^3 \langle \bar{q} g_s \sigma G q \rangle}{64\pi^2 T^4} \int_{m_c^2}^{s_{D^*}^0} du u \left(1 - \frac{m_c^2}{u} \right)^2 \exp\left(-\frac{u+m_c^2}{T^2}\right) - \frac{m_c \langle \bar{q} g_s \sigma G q \rangle}{192\pi^2 T^2} \left(2 - \frac{m_c^2}{T^2} \right) \int_{m_c^2}^{s_{D_1}^0} ds \left(1 - \frac{m_c^2}{s} \right)^2 \\ &\times (2s+m_c^2) \exp\left(-\frac{s+m_c^2}{T^2}\right) + \frac{m_c \langle \bar{q} g_s \sigma G q \rangle}{192\pi^2} \int_{m_c^2}^{s_{D^*}^0} du \left(3 - \frac{2m_c^2}{u} \right) \exp\left(-\frac{u+m_c^2}{T^2}\right) \\ &- \frac{m_c^3 \langle \bar{q} g_s \sigma G q \rangle}{192\pi^2} \int_{m_c^2}^{s_{D_1}^0} ds \frac{1}{s} \exp\left(-\frac{s+m_c^2}{T^2}\right), \end{aligned} \quad (\text{C4})$$

$$\begin{aligned} \Pi_{\eta_c \omega PA}^{\text{QCD}}(T^2) = & -\frac{m_c \langle \bar{q}q \rangle}{2\sqrt{2}\pi^2} \int_{4m_c^2}^{s_{\eta_c}^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \exp\left(-\frac{s}{T^2}\right) + \frac{m_c \langle \bar{q}g_s \sigma G q \rangle}{6\sqrt{2}\pi^2 T^2} \int_{4m_c^2}^{s_{\eta_c}^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \exp\left(-\frac{s}{T^2}\right) \\ & - \frac{m_c \langle \bar{q}g_s \sigma G q \rangle}{24\sqrt{2}\pi^2} \int_{4m_c^2}^{s_{\eta_c}^0} ds \frac{1}{\sqrt{s(s-4m_c^2)}} \exp\left(-\frac{s}{T^2}\right), \end{aligned} \quad (\text{C5})$$

$$\Pi_{J/\psi \omega PA}^{\text{QCD}}(T^2) = 0, \quad (\text{C6})$$

$$\Pi_{\chi_{c0} \omega PA}^{\text{QCD}}(T^2) = \frac{1}{32\sqrt{2}\pi^4} \int_{4m_c^2}^{s_{\chi_{c0}}^0} ds \int_0^{s_{\chi_{c0}}^0} du \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} u (s-4m_c^2) \exp\left(-\frac{s+u}{T^2}\right), \quad (\text{C7})$$

$$\begin{aligned} \Pi_{\chi_{c1} \omega PA}^{\text{QCD}}(T^2) = & -\frac{\langle \bar{q}q \rangle}{12\sqrt{2}\pi^2} \int_{4m_c^2}^{s_{\chi_{c1}}^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \left(1 + \frac{2m_c^2}{s}\right) \exp\left(-\frac{s}{T^2}\right) \\ & + \frac{\langle \bar{q}g_s \sigma G q \rangle}{36\sqrt{2}\pi^2 T^2} \int_{4m_c^2}^{s_{\chi_{c1}}^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \left(1 + \frac{2m_c^2}{s}\right) \exp\left(-\frac{s}{T^2}\right), \end{aligned} \quad (\text{C8})$$

$$\Pi_{J/\psi f_0(500) PA}^{\text{QCD}}(T^2) = \frac{1}{32\sqrt{2}\pi^4} \int_{4m_c^2}^{s_{J/\psi}^0} ds \int_0^{s_{f_0(500)}^0} du \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} u (s+2m_c^2) \exp\left(-\frac{s+u}{T^2}\right), \quad (\text{C9})$$

C.2. The QCD side for the current $\tilde{J}_\mu^{\text{AV}}$

$$\begin{aligned} \Pi_{D_s D_s \tilde{A} V}^{\text{QCD}}(T^2) = & \frac{9}{64\sqrt{2}\pi^4} \int_{m_c^2}^{s_{D_s}^0} ds \int_{m_c^2}^{s_{D_s}^0} du \left(1 - \frac{m_c^2}{s}\right) \left(1 - \frac{m_c^2}{u}\right) \exp\left(-\frac{s+u}{T^2}\right) \\ & - \frac{m_c (s-m_c^2)(u-m_c^2) + m_s (su+m_c^2(s+u)-3m_c^4)}{8\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_s}^0} ds \left(1 - \frac{m_c^2}{s}\right) \\ & \times \frac{m_s (s+m_c^2) + m_c s - m_c^3}{s} \exp\left(-\frac{s+m_c^2}{T^2}\right) - \frac{3\langle \bar{s}s \rangle}{8\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_s}^0} du \left(1 - \frac{m_c^2}{u}\right) (2m_s m_c + u - m_c^2) \exp\left(-\frac{u+m_c^2}{T^2}\right) \\ & + \frac{3m_s m_c \langle \bar{s}s \rangle}{16\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_s}^0} ds \left(1 - \frac{m_c^2}{s}\right)^2 \left(1 - \frac{s-m_c^2}{T^2}\right) \exp\left(-\frac{s+m_c^2}{T^2}\right) + \frac{3m_c^3 \langle \bar{s}g_s \sigma G s \rangle}{32\sqrt{2}\pi^2 T^4} \int_{m_c^2}^{s_{D_s}^0} ds \left(1 - \frac{m_c^2}{s}\right) \\ & \times \frac{m_s (s+m_c^2) + m_c s - m_c^3}{s} \exp\left(-\frac{s+m_c^2}{T^2}\right) + \frac{3m_c^2 \langle \bar{s}g_s \sigma G s \rangle}{32\sqrt{2}\pi^2 T^4} \int_{m_c^2}^{s_{D_s}^0} du \left(1 - \frac{m_c^2}{u}\right) (2m_s m_c + u - m_c^2) \\ & \times \exp\left(-\frac{u+m_c^2}{T^2}\right) + \frac{m_s m_c^3 \langle \bar{s}g_s \sigma G s \rangle}{32\sqrt{2}\pi^2 T^6} \int_{m_c^2}^{s_{D_s}^0} ds \left(1 - \frac{m_c^2}{s}\right)^2 (s-m_c^2) \exp\left(-\frac{s+m_c^2}{T^2}\right) \\ & - \frac{3m_c \langle \bar{s}g_s \sigma G s \rangle}{16\sqrt{2}\pi^2 T^2} \int_{m_c^2}^{s_{D_s}^0} ds \left(1 - \frac{m_c^2}{s}\right) \frac{m_s (s+m_c^2) + m_c s - m_c^3}{s} \exp\left(-\frac{s+m_c^2}{T^2}\right) + \frac{m_c^2 \langle \bar{s}g_s \sigma G s \rangle}{32\sqrt{2}\pi^2} \\ & \times \int_{m_c^2}^{s_{D_s}^0} ds \frac{m_s m_c + s - m_c^2}{s^2} \exp\left(-\frac{s+m_c^2}{T^2}\right) + \frac{\langle \bar{s}g_s \sigma G s \rangle}{192\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_s}^0} du \frac{6m_s m_c u + 3u^2 - 4m_c^2 u + m_c^4}{u^2} \exp\left(-\frac{u+m_c^2}{T^2}\right) \\ & + \frac{m_c \langle \bar{s}g_s \sigma G s \rangle}{32\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_s}^0} ds \frac{m_s (s^2+m_c^4) + m_c^3 s - m_c^5}{s^2 (s-m_c^2)} \exp\left(-\frac{s+m_c^2}{T^2}\right) + \frac{\langle \bar{s}g_s \sigma G s \rangle}{192\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_s}^0} du \exp\left(-\frac{u+m_c^2}{T^2}\right) \\ & \frac{6m_s m_c^3 u + 6m_s m_c u^2 + m_c^6 - m_c^4 u - 3m_c^2 u^2 + 3u^3}{u^2 (u-m_c^2)}, \end{aligned} \quad (\text{C10})$$

$$\begin{aligned}
\Pi_{D_s^* D_s \tilde{A} V}^{\text{QCD}}(T^2) = & -\frac{m_c \langle \bar{s} g_s \sigma G s \rangle}{96 \sqrt{2} \pi^2} \int_{m_c^2}^{s_0^0} du \frac{2m_s m_c + 3m_c^2 - u}{u^2} \exp\left(-\frac{u+m_c^2}{T^2}\right) - \frac{m_c \langle \bar{s} g_s \sigma G s \rangle}{96 \sqrt{2} \pi^2} \int_{m_c^2}^{s_0^0} ds \frac{2m_s m_c + 3m_c^2 + s}{s^2} \\
& \times \exp\left(-\frac{s+m_c^2}{T^2}\right) - \frac{\langle \bar{s} g_s \sigma G s \rangle}{96 \sqrt{2} \pi^2} \int_{m_c^2}^{s_0^0} du \frac{5m_s (u^2 + m_c^4)}{u^2 (u-m_c^2)} \exp\left(-\frac{u+m_c^2}{T^2}\right) \\
& - \frac{\langle \bar{s} g_s \sigma G s \rangle}{96 \sqrt{2} \pi^2} \int_{m_c^2}^{s_0^0} ds \frac{5m_s (s^2 + m_c^4)}{s^2 (s-m_c^2)} \exp\left(-\frac{s+m_c^2}{T^2}\right), \quad (\text{C11})
\end{aligned}$$

$$\begin{aligned}
\Pi_{D_s^* D_s \tilde{A} V}^{\text{QCD}}(T^2) = & -\frac{1}{64 \sqrt{2} \pi^4} \int_{m_c^2}^{s_0^0} ds \int_{m_c^2}^{s_0^0} du \left(1 - \frac{m_c^2}{s}\right) \left(1 - \frac{m_c^2}{u}\right) \exp\left(-\frac{s+u}{T^2}\right) \\
& \frac{m_s (m_c^6 + m_c^4 (s+7u) - m_c^2 u (5s+2u) - 2su^2) - m_c (s-m_c^2) (u-m_c^2) (2u+m_c^2)}{su} \\
& - \frac{m_c \langle \bar{s} s \rangle}{8 \sqrt{2} \pi^2} \int_{m_c^2}^{s_0^0} ds \left(1 - \frac{m_c^2}{s}\right) \frac{m_s (s+m_c^2) + m_c (s-m_c^2)}{s} \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& - \frac{\langle \bar{s} s \rangle}{24 \sqrt{2} \pi^2} \int_{m_c^2}^{s_0^0} du \left(1 - \frac{m_c^2}{u}\right) \frac{6m_s m_c u + (u-m_c^2) (2u+m_c^2)}{u} \exp\left(-\frac{u+m_c^2}{T^2}\right) \\
& - \frac{3m_s m_c \langle \bar{s} s \rangle}{8 \sqrt{2} \pi^2 T^2} \int_{m_c^2}^{s_0^0} ds \left(1 - \frac{m_c^2}{s}\right)^2 s \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& + \frac{m_c^3 \langle \bar{s} g_s \sigma G s \rangle}{32 \sqrt{2} \pi^2 T^4} \int_{m_c^2}^{s_0^0} ds \left(1 - \frac{m_c^2}{s}\right) \frac{m_s (s+m_c^2) + m_c (s-m_c^2)}{s} \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& + \frac{\langle \bar{s} g_s \sigma G s \rangle}{288 \sqrt{2} \pi^2 T^2} \left(4 + \frac{3m_c^2}{T^2}\right) \int_{m_c^2}^{s_0^0} du \left(1 - \frac{m_c^2}{u}\right) \exp\left(-\frac{u+m_c^2}{T^2}\right) \frac{6m_s m_c u + (u-m_c^2) (2u+m_c^2)}{u} \\
& + \frac{m_s m_c \langle \bar{s} g_s \sigma G s \rangle}{96 \sqrt{2} \pi^2 T^2} \left(1 + \frac{m_c^2}{T^2} - \frac{m_c^4}{T^4}\right) \int_{m_c^2}^{s_0^0} ds \left(1 - \frac{m_c^2}{s}\right)^2 \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& + \frac{m_s m_c^3 \langle \bar{s} g_s \sigma G s \rangle}{288 \sqrt{2} \pi^2 T^6} \int_{m_c^2}^{s_0^0} du \left(1 - \frac{m_c^2}{u}\right)^2 (2u+m_c^2) \exp\left(-\frac{u+m_c^2}{T^2}\right) \\
& + \frac{m_c^2 \langle \bar{s} g_s \sigma G s \rangle}{96 \sqrt{2} \pi^2} \int_{m_c^2}^{s_0^0} ds \frac{m_s m_c + s + 4m_c^2}{s^2} \exp\left(-\frac{s+m_c^2}{T^2}\right) + \frac{\langle \bar{s} g_s \sigma G s \rangle}{96 \sqrt{2} \pi^2} \int_{m_c^2}^{s_0^0} du \frac{m_s m_c + m_c^2}{u} \exp\left(-\frac{u+m_c^2}{T^2}\right) \\
& + \frac{5m_c \langle \bar{s} g_s \sigma G s \rangle}{96 \sqrt{2} \pi^2} \int_{m_c^2}^{s_0^0} ds \frac{m_s (s^2 + m_c^4)}{s^2 (s-m_c^2)} \exp\left(-\frac{s+m_c^2}{T^2}\right) + \frac{\langle \bar{s} g_s \sigma G s \rangle}{96 \sqrt{2} \pi^2} \int_{m_c^2}^{s_0^0} du \frac{5m_s m_c (u+m_c^2)}{u (u-m_c^2)} \exp\left(-\frac{u+m_c^2}{T^2}\right), \quad (\text{C12})
\end{aligned}$$

$$\Pi_{J/\psi \omega \tilde{A} V}^{\text{QCD}}(T^2) = 0, \quad (\text{C13})$$

$$\begin{aligned}
\Pi_{\chi_{c0} \omega \tilde{A} V}^{\text{QCD}}(T^2) = & \frac{3}{32 \sqrt{2} \pi^4} \int_{4m_c^2}^{s_0^0} ds \int_0^{s_0^0} du \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} u (s-4m_c^2) \exp\left(-\frac{s+u}{T^2}\right) \\
& - \frac{3m_s \langle \bar{s} s \rangle}{4 \sqrt{2} \pi^2} \int_{4m_c^2}^{s_0^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} (s-4m_c^2) \exp\left(-\frac{s}{T^2}\right) \\
& + \frac{m_s \langle \bar{s} g_s \sigma G s \rangle}{48 \sqrt{2} \pi^2} \int_{4m_c^2}^{s_0^0} ds \frac{3s-16m_c^2}{\sqrt{s(s-4m_c^2)}} \exp\left(-\frac{s}{T^2}\right), \quad (\text{C14})
\end{aligned}$$

$$\begin{aligned}
\Pi_{D_{s0}D_s^*AV}^{\text{QCD}}(T^2) = & \frac{3m_c^2}{64\sqrt{2}\pi^4} \int_{m_c^2}^{s_{D_{s0}}^0} ds \int_{m_c^2}^{s_{D_s^*}^0} du \left(1 - \frac{m_c^2}{s}\right) \left(1 - \frac{m_c^2}{u}\right) \exp\left(-\frac{s+u}{T^2}\right) \frac{2m_s m_c (s-u) + (s-m_c^2)(u-m_c^2)}{su} \\
& + \frac{\langle \bar{s}s \rangle}{8\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_{s0}}^0} ds \left(1 - \frac{m_c^2}{s}\right) \frac{m_s(s+m_c^2) - m_c(s-m_c^2)}{s} \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& + \frac{\langle \bar{s}s \rangle}{8\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_s^*}^0} du \left(1 - \frac{m_c^2}{u}\right) \frac{m_s(u+m_c^2) + m_c(u-m_c^2)}{u} \exp\left(-\frac{u+m_c^2}{T^2}\right) \\
& + \frac{m_s m_c^2 \langle \bar{s}s \rangle}{16\sqrt{2}\pi^2 T^2} \int_{m_c^2}^{s_{D_{s0}}^0} ds \left(1 - \frac{m_c^2}{s}\right)^2 \exp\left(-\frac{s+m_c^2}{T^2}\right) - \frac{m_s m_c^2 \langle \bar{s}s \rangle}{16\sqrt{2}\pi^2 T^2} \int_{m_c^2}^{s_{D_s^*}^0} du \left(1 - \frac{m_c^2}{u}\right)^2 \exp\left(-\frac{u+m_c^2}{T^2}\right) \\
& - \frac{\langle \bar{s}g_s \sigma G s \rangle}{96\sqrt{2}\pi^2 T^2} \left(4 + \frac{3m_c^2}{T^2}\right) \int_{m_c^2}^{s_{D_{s0}}^0} ds \left(1 - \frac{m_c^2}{s}\right) \exp\left(-\frac{s+m_c^2}{T^2}\right) \frac{m_s(s+m_c^2) - m_c(s-m_c^2)}{s} \\
& - \frac{m_c^2 \langle \bar{s}g_s \sigma G s \rangle}{32\sqrt{2}\pi^2 T^4} \int_{m_c^2}^{s_{D_s^*}^0} du \left(1 - \frac{m_c^2}{u}\right) \frac{m_s(u+m_c^2) + m_c(u-m_c^2)}{u} \exp\left(-\frac{u+m_c^2}{T^2}\right) \\
& - \frac{m_s m_c^4 \langle \bar{s}g_s \sigma G s \rangle}{96\sqrt{2}\pi^2 T^6} \int_{m_c^2}^{s_{D_{s0}}^0} ds \left(1 - \frac{m_c^2}{s}\right)^2 \exp\left(-\frac{s+m_c^2}{T^2}\right) + \frac{m_s m_c^4 \langle \bar{s}g_s \sigma G s \rangle}{96\sqrt{2}\pi^2 T^6} \int_{m_c^2}^{s_{D_s^*}^0} du \left(1 - \frac{m_c^2}{u}\right)^2 \\
& \times \exp\left(-\frac{u+m_c^2}{T^2}\right) - \frac{m_c \langle \bar{s}g_s \sigma G s \rangle}{96\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_{s0}}^0} ds \frac{3m_s m_c - s - 2m_c^2}{s^2} \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& - \frac{m_c \langle \bar{s}g_s \sigma G s \rangle}{96\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_s^*}^0} du \frac{3m_s m_c + u + 2m_c^2}{u^2} \exp\left(-\frac{u+m_c^2}{T^2}\right) - \frac{\langle \bar{s}g_s \sigma G s \rangle}{96\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_{s0}}^0} ds \frac{5m_s(s^2+m_c^4)}{s^2(s-m_c^2)} \\
& \times \exp\left(-\frac{s+m_c^2}{T^2}\right) - \frac{\langle \bar{s}g_s \sigma G s \rangle}{96\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_s^*}^0} du \frac{5m_s(u^2+m_c^4)}{u^2(u-m_c^2)} \exp\left(-\frac{u+m_c^2}{T^2}\right),
\end{aligned} \tag{C15}$$

$$\begin{aligned}
\Pi_{\chi_{c1}\omega AV}^{\text{QCD}}(T^2) = & \frac{m_s}{32\sqrt{2}\pi^4} \int_{4m_c^2}^{s_{\chi_{c1}}^0} ds \int_0^{s_\phi^0} du \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \left(1 + \frac{2m_c^2}{s}\right) \exp\left(-\frac{s+u}{T^2}\right) - \frac{\langle \bar{s}s \rangle}{12\sqrt{2}\pi^2} \int_{4m_c^2}^{s_{\chi_{c1}}^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \\
& \times \left(1 + \frac{2m_c^2}{s}\right) \exp\left(-\frac{s}{T^2}\right) + \frac{\langle \bar{s}g_s \sigma G s \rangle}{36\sqrt{2}\pi^2 T^2} \int_{4m_c^2}^{s_{\chi_{c1}}^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \left(1 + \frac{2m_c^2}{s}\right) \exp\left(-\frac{s}{T^2}\right),
\end{aligned} \tag{C16}$$

$$\begin{aligned}
\Pi_{D_{s1}D_s AV}^{\text{QCD}}(T^2) = & \frac{3}{64\sqrt{2}\pi^4} \int_{m_c^2}^{s_{D_{s1}}^0} ds \int_{m_c^2}^{s_{D_s}^0} du \left(1 - \frac{m_c^2}{s}\right) \left(1 - \frac{m_c^2}{u}\right) \exp\left(-\frac{s+u}{T^2}\right) \\
& \frac{(s-m_c^2)(2s+m_c^2)(u-m_c^2) - 6m_s m_c (m_c^4 - 2m_c^2 s + m_c s(3u-2s))}{s} \\
& + \frac{m_c \langle \bar{s}s \rangle}{8\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_{s1}}^0} ds \left(1 - \frac{m_c^2}{s}\right) \frac{6m_s m_c s - (s-m_c^2)(2s+m_c^2)}{s} \exp\left(-\frac{s+m_c^2}{T^2}\right) + \frac{3m_c \langle \bar{s}s \rangle}{8\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_s}^0} du \left(1 - \frac{m_c^2}{u}\right) \\
& \times (2m_s m_c + u - m_c^2) \exp\left(-\frac{u+m_c^2}{T^2}\right) + \frac{m_s \langle \bar{s}s \rangle}{16\sqrt{2}\pi^2} \left(1 + \frac{m_c^2}{T^2}\right) \int_{m_c^2}^{s_{D_{s1}}^0} ds \left(1 - \frac{m_c^2}{s}\right)^2 (2s+m_c^2) \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& - \frac{3m_s m_c^2 \langle \bar{s}s \rangle}{16\sqrt{2}\pi^2 T^2} \int_{m_c^2}^{s_{D_s}^0} du \left(1 - \frac{m_c^2}{u}\right)^2 u \exp\left(-\frac{u+m_c^2}{T^2}\right) + \frac{m_c \langle \bar{s}g_s \sigma G s \rangle}{32\sqrt{2}\pi^2 T^2} \left(2 - \frac{m_c^2}{T^2}\right) \\
& \times \int_{m_c^2}^{s_{D_{s1}}^0} ds \left(1 - \frac{m_c^2}{s}\right) \exp\left(-\frac{s+m_c^2}{T^2}\right) \frac{6m_s m_c s - (s-m_c^2)(2s+m_c^2)}{s} - \frac{3m_c^3 \langle \bar{s}g_s \sigma G s \rangle}{32\sqrt{2}\pi^2 T^4}
\end{aligned}$$

$$\begin{aligned}
& \times \int_{m_c^2}^{s_{D_s}^0} du \left(1 - \frac{m_c^2}{u}\right)^2 (2m_s m_c + u - m_c^2) \exp\left(-\frac{u+m_c^2}{T^2}\right) - \frac{m_s m_c^4 \langle \bar{s} g_s \sigma G s \rangle}{96 \sqrt{2} \pi^2 T^6} \int_{m_c^2}^{s_{D_{s1}}^0} ds \left(1 - \frac{m_c^2}{s}\right)^2 \\
& \times (2s + m_c^2) \exp\left(-\frac{s+m_c^2}{T^2}\right) + \frac{m_s \langle \bar{s} g_s \sigma G s \rangle}{32 \sqrt{2} \pi^2 T^2} \left(-1 - \frac{m_c^2}{T^2} + \frac{m_c^4}{T^4}\right) \int_{m_c^2}^{s_{D_s}^0} du \left(1 - \frac{m_c^2}{u}\right)^2 u \exp\left(-\frac{u+m_c^2}{T^2}\right) \\
& + \frac{m_c^2 \langle \bar{s} g_s \sigma G s \rangle}{96 \sqrt{2} \pi^2} \int_{m_c^2}^{s_{D_{s1}}^0} ds \frac{3m_s s - m_c (s - 2m_c^2)}{s^2} \exp\left(-\frac{s+m_c^2}{T^2}\right) - \frac{m_c \langle \bar{s} g_s \sigma G s \rangle}{192 \sqrt{2} \pi^2} \\
& \times \int_{m_c^2}^{s_{D_s}^0} du \frac{6m_s m_c u + 2u (3u - 2m_c^2)}{u^2} \exp\left(-\frac{u+m_c^2}{T^2}\right) - \frac{m_c^2 \langle \bar{s} g_s \sigma G s \rangle}{96 \sqrt{2} \pi^2} \int_{m_c^2}^{s_{D_{s1}}^0} ds \frac{3m_s (s + m_c^2)}{s (s - m_c^2)} \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& - \frac{m_c \langle \bar{s} g_s \sigma G s \rangle}{192 \sqrt{2} \pi^2} \int_{m_c^2}^{s_{D_s}^0} du \frac{6m_s m_c (u + m_c^2)}{u (u - m_c^2)} \exp\left(-\frac{u+m_c^2}{T^2}\right),
\end{aligned} \tag{C17}$$

$$\begin{aligned}
\Pi_{\eta_c \phi_{AV}}^{\text{QCD}}(T^2) &= \frac{3m_s m_c}{16 \sqrt{2} \pi^4} \int_{4m_c^2}^{s_{\eta_c}^0} ds \int_0^{s_{\phi}^0} du \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \exp\left(-\frac{s+u}{T^2}\right) - \frac{m_c \langle \bar{s} s \rangle}{\sqrt{2} \pi^2} \int_{4m_c^2}^{s_{\eta_c}^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \exp\left(-\frac{s}{T^2}\right) \\
&+ \frac{m_c \langle \bar{s} g_s \sigma G s \rangle}{3 \sqrt{2} \pi^2 T^2} \int_{4m_c^2}^{s_{\eta_c}^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \exp\left(-\frac{s}{T^2}\right) + \frac{m_c \langle \bar{s} g_s \sigma G s \rangle}{4 \sqrt{2} \pi^2} \int_{4m_c^2}^{s_{\eta_c}^0} ds \frac{1}{\sqrt{s(s-4m_c^2)}} \exp\left(-\frac{s}{T^2}\right),
\end{aligned} \tag{C18}$$

$$\begin{aligned}
\Pi_{J/\psi f_0(980)AV}^{\text{QCD}}(T^2) &= \frac{3}{32 \sqrt{2} \pi^4} \int_{4m_c^2}^{s_{J/\psi}^0} ds \int_0^{s_{f_0(980)}^0} du \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} u (s + 2m_c^2) \exp\left(-\frac{s+u}{T^2}\right) + \frac{m_s \langle \bar{s} s \rangle}{2 \sqrt{2} \pi^2} \int_{4m_c^2}^{s_{J/\psi}^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \\
&\times (s + 2m_c^2) \exp\left(-\frac{s}{T^2}\right) - \frac{m_s m_c^2 \langle \bar{s} g_s \sigma G s \rangle}{6 \sqrt{2} \pi^2} \int_{4m_c^2}^{s_{J/\psi}^0} ds \frac{1}{\sqrt{s(s-4m_c^2)}} \exp\left(-\frac{s}{T^2}\right) \\
&+ \frac{m_s \langle \bar{s} g_s \sigma G s \rangle}{4 \sqrt{2} \pi^2 T^2} \int_{4m_c^2}^{s_{J/\psi}^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} (s + 2m_c^2) \exp\left(-\frac{s}{T^2}\right),
\end{aligned} \tag{C19}$$

C.3. The QCD side for the current $J_{\mu\nu}^{AA}$

$$\Pi_{DDAA}^{\text{QCD}}(T^2) = -\frac{m_c^2 \langle \bar{q} g_s \sigma G q \rangle}{48 \pi^2} \int_{m_c^2}^{s_D^0} ds \frac{1}{s^2} \exp\left(-\frac{s+m_c^2}{T^2}\right), \tag{C20}$$

$$\Pi_{D^* DAA}^{\text{QCD}}(T^2) = -\frac{m_c^2 \langle \bar{q} g_s \sigma G q \rangle}{96 \pi^2} \int_{m_c^2}^{s_{D^*}^0} du \frac{1}{u^2} \exp\left(-\frac{u+m_c^2}{T^2}\right), \tag{C21}$$

$$\Pi_{D^* D^* AA}^{\text{QCD}}(T^2) = 0, \tag{C22}$$

$$\Pi_{D_0 D^* AA}^{\text{QCD}}(T^2) = \frac{3m_c}{128 \pi^4} \int_{m_c^2}^{s_{D_0}^0} ds \int_{m_c^2}^{s_{D^*}^0} du \left(1 - \frac{m_c^2}{s}\right)^2 \left(1 - \frac{m_c^2}{u}\right)^2 s \exp\left(-\frac{s+u}{T^2}\right) - \frac{\langle \bar{q} q \rangle}{16 \pi^2} \int_{m_c^2}^{s_{D_0}^0} ds \left(1 - \frac{m_c^2}{s}\right)^2 s \exp\left(-\frac{s+m_c^2}{T^2}\right)$$

$$\begin{aligned}
& + \frac{m_c^2 \langle \bar{q}q \rangle}{16\pi^2} \int_{m_c^2}^{s_{D^*}^0} du \left(1 - \frac{m_c^2}{u}\right)^2 \exp\left(-\frac{u+m_c^2}{T^2}\right) + \frac{\langle \bar{q}g_s \sigma G q \rangle}{192\pi^2 T^2} \left(4 + \frac{3m_c^2}{T^2}\right) \int_{m_c^2}^{s_{D_0}^0} ds \left(1 - \frac{m_c^2}{s}\right)^2 s \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& + \frac{m_c^2 \langle \bar{q}g_s \sigma G q \rangle}{64\pi^2 T^2} \left(2 - \frac{m_c^2}{T^2}\right) \int_{m_c^2}^{s_{D^*}^0} du \left(1 - \frac{m_c^2}{u}\right)^2 \exp\left(-\frac{u+m_c^2}{T^2}\right) + \frac{m_c^2 \langle \bar{q}g_s \sigma G q \rangle}{192\pi^2} \int_{m_c^2}^{s_{D_0}^0} ds \frac{1}{s} \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& + \frac{m_c^4 \langle \bar{q}g_s \sigma G q \rangle}{192\pi^2} \int_{m_c^2}^{s_{D^*}^0} du \frac{1}{u^2} \exp\left(-\frac{u+m_c^2}{T^2}\right), \tag{C23}
\end{aligned}$$

$$\begin{aligned}
\Pi_{D_1 D A A}^{\text{QCD}}(T^2) = & \frac{3m_c}{128\pi^4} \int_{m_c^2}^{s_{D_1}^0} ds \int_{m_c^2}^{s_D^0} du \left(1 - \frac{m_c^2}{s}\right)^2 \left(1 - \frac{m_c^2}{u}\right)^2 u \exp\left(-\frac{s+u}{T^2}\right) - \frac{m_c^2 \langle \bar{q}q \rangle}{16\pi^2} \int_{m_c^2}^{s_{D_1}^0} ds \left(1 - \frac{m_c^2}{s}\right)^2 \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& + \frac{\langle \bar{q}q \rangle}{16\pi^2} \int_{m_c^2}^{s_D^0} du \left(1 - \frac{m_c^2}{u}\right)^2 u \exp\left(-\frac{u+m_c^2}{T^2}\right) - \frac{m_c^2 \langle \bar{q}g_s \sigma G q \rangle}{64\pi^2 T^2} \left(2 - \frac{m_c^2}{T^2}\right) \int_{m_c^2}^{s_{D_1}^0} ds \left(1 - \frac{m_c^2}{s}\right)^2 \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& - \frac{\langle \bar{q}g_s \sigma G q \rangle}{192\pi^2 T^2} \left(4 + \frac{3m_c^2}{T^2}\right) \int_{m_c^2}^{s_D^0} du \left(1 - \frac{m_c^2}{u}\right)^2 u \exp\left(-\frac{u+m_c^2}{T^2}\right) - \frac{m_c^2 \langle \bar{q}g_s \sigma G q \rangle}{192\pi^2} \int_{m_c^2}^{s_D^0} du \frac{1}{u} \exp\left(-\frac{u+m_c^2}{T^2}\right) \\
& - \frac{m_c^4 \langle \bar{q}g_s \sigma G q \rangle}{192\pi^2} \int_{m_c^2}^{s_{D_1}^0} ds \frac{1}{s^2} \exp\left(-\frac{s+m_c^2}{T^2}\right), \tag{C24}
\end{aligned}$$

$$\Pi_{\eta_c \omega A A}^{\text{QCD}}(T^2) = -\frac{m_c^2 \langle \bar{q}g_s \sigma G q \rangle}{12\sqrt{2}\pi^2} \int_{4m_c^2}^{s_{\eta_c}^0} ds \frac{1}{\sqrt{s(s-4m_c^2)}} \exp\left(-\frac{s}{T^2}\right), \tag{C25}$$

$$\Pi_{J/\psi \omega A A}^{\text{QCD}}(T^2) = 0, \tag{C26}$$

$$\begin{aligned}
\Pi_{\chi_{c0} \omega A A}^{\text{QCD}}(T^2) = & -\frac{\langle \bar{q}q \rangle}{4\sqrt{2}\pi^2} \int_{4m_c^2}^{s_{\chi_{c0}}^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} (s-4m_c^2) \exp\left(-\frac{s}{T^2}\right) + \frac{\langle \bar{q}g_s \sigma G q \rangle}{12\sqrt{2}\pi^2 T^2} \int_{4m_c^2}^{s_{\chi_{c0}}^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \\
& \times (s-4m_c^2) \exp\left(-\frac{s}{T^2}\right) + \frac{m_c^2 \langle \bar{q}g_s \sigma G q \rangle}{24\sqrt{2}\pi^2} \int_{4m_c^2}^{s_{\chi_{c0}}^0} ds \frac{1}{\sqrt{s(s-4m_c^2)}} \exp\left(-\frac{s}{T^2}\right), \tag{C27}
\end{aligned}$$

$$\Pi_{\chi_{c1} \omega A A}^{\text{QCD}}(T^2) = \frac{1}{24\sqrt{2}\pi^4} \int_{4m_c^2}^{s_{\chi_{c1}}^0} ds \int_0^{s_\omega^0} du \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} u (7s-40m_c^2) \exp\left(-\frac{s+u}{T^2}\right), \tag{C28}$$

$$\Pi_{J/\psi f_0(500) A A}^{\text{QCD}}(T^2) = \frac{3m_c}{32\sqrt{2}\pi^4} \int_{4m_c^2}^{s_{J/\psi}^0} ds \int_0^{s_{f_0(500)}^0} du \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} u \exp\left(-\frac{s+u}{T^2}\right), \tag{C29}$$

C.4. The QCD side for the current $J_{\mu\nu}^{\widetilde{V}}$

$$\Pi_{D_s D_s \widetilde{V}}^{\text{QCD}}(T^2) = \frac{3m_c}{128\sqrt{2}\pi^4} \int_{m_c^2}^{s_{D_s}^0} ds \int_{m_c^2}^{s_{D_s}^0} du \left(1 - \frac{m_c^2}{s}\right) \left(1 - \frac{m_c^2}{u}\right) \exp\left(-\frac{s+u}{T^2}\right) \frac{2m_s(su-m_c^4) + m_c(s-m_c^2)(u-m_c^2)}{su}$$

$$\begin{aligned}
& -\frac{\langle \bar{s}s \rangle}{8\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_s^0}} ds \left(1 - \frac{m_c^2}{s}\right) \frac{m_s(s+m_c^2) + m_c(s-m_c^2)}{s} \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& + \frac{m_c^2 \langle \bar{s}g_s \sigma G s \rangle}{32\sqrt{2}\pi^2 T^4} \int_{m_c^2}^{s_{D_s^0}} ds \left(1 - \frac{m_c^2}{s}\right) \frac{m_s(s+m_c^2) + m_c(s-m_c^2)}{s} \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& - \frac{m_c \langle \bar{s}g_s \sigma G s \rangle}{96\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_s^0}} ds \frac{m_s m_c + 2s - m_c^2}{s^2} \exp\left(-\frac{s+m_c^2}{T^2}\right), \tag{C30}
\end{aligned}$$

$$\Pi_{D_s^* D_s^* S \tilde{V}}^{\text{QCD}}(T^2) = -\frac{m_c^4 \langle \bar{s}g_s \sigma G s \rangle}{48\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_s^*}^0} ds \frac{1}{s^3} \exp\left(-\frac{s+m_c^2}{T^2}\right), \tag{C31}$$

$$\begin{aligned}
\Pi_{D_s^* D_s^* S \tilde{V}}^{\text{QCD}}(T^2) = & \frac{3m_c}{256\sqrt{2}\pi^4} \int_{m_c^2}^{s_{D_s^*}^0} ds \int_{m_c^2}^{s_{D_s^*}^0} du \left(1 - \frac{m_c^2}{s}\right)^2 \frac{m_s(u^2 - m_c^4) + m_c(u - m_c^2)^2}{u^2} \exp\left(-\frac{s+u}{T^2}\right) \\
& - \frac{\langle \bar{s}s \rangle}{32\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_s^*}^0} ds \frac{2m_s(s^2 - m_c^4) + m_c(s - m_c^2)^2}{s^2} \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& + \frac{\langle \bar{s}g_s \sigma G s \rangle}{128\sqrt{2}\pi^2 T^2} \left(1 + \frac{m_c^2}{T^2}\right) \int_{m_c^2}^{s_{D_s^*}^0} ds \frac{2m_s(s^2 - m_c^4) + m_c(s - m_c^2)^2}{s^2} \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& + \frac{\langle \bar{s}g_s \sigma G s \rangle}{384\sqrt{2}\pi^2 T^2} \int_{m_c^2}^{s_{D_s^*}^0} ds \frac{m_s(s^2 - m_c^4) + 2m_c(s - m_c^2)^2}{s^2} \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& - \frac{m_c \langle \bar{s}g_s \sigma G s \rangle}{384\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_s^*}^0} ds \frac{1}{s} \left(1 - \frac{m_c^2}{s}\right) \exp\left(-\frac{s+m_c^2}{T^2}\right) + \frac{m_s m_c^2 \langle \bar{s}g_s \sigma G s \rangle}{384\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_s^*}^0} du \frac{1}{u^2} \exp\left(-\frac{u+m_c^2}{T^2}\right) \\
& - \frac{\langle \bar{s}g_s \sigma G s \rangle}{768\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_s^*}^0} ds \frac{2m_s(s^2 + m_c^4) + 3m_c^3(s - m_c^2)}{s^2(s - m_c^2)} \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& - \frac{m_c \langle \bar{s}g_s \sigma G s \rangle}{1536\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_s^*}^0} du \frac{2m_s m_c(3u - m_c^2) + 2u^2 - m_c^2 u + m_c^4}{u^2(u - m_c^2)} \exp\left(-\frac{u+m_c^2}{T^2}\right), \tag{C32}
\end{aligned}$$

$$\begin{aligned}
\Pi_{D_{s0} D_s^* S \tilde{V}}^{\text{QCD}}(T^2) = & \frac{1}{64\sqrt{2}\pi^4} \int_{m_c^2}^{s_{D_{s0}}^0} ds \int_{m_c^2}^{s_{D_s^*}^0} du \left(1 - \frac{m_c^2}{s}\right) \left(1 - \frac{m_c^2}{u}\right) \exp\left(-\frac{s+u}{T^2}\right) \\
& \frac{m_s[m_c^6 + m_c^4(s-5u) + m_c^2 u(7s-2u) - 2su^2] + m_c(2u+m_c^2)(s-m_c^2)(u-m_c^2)}{su} \\
& + \frac{m_c \langle \bar{s}s \rangle}{8\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_{s0}}^0} ds \left(1 - \frac{m_c^2}{s}\right) \frac{m_s(s+m_c^2) - m_c(s-m_c^2)}{s} \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& - \frac{\langle \bar{s}s \rangle}{24\sqrt{2}\pi^2} \int_{m_c^2}^{s_{D_s^*}^0} du \left(1 - \frac{m_c^2}{u}\right) \frac{6m_s m_c u + (2u+m_c^2)(u-m_c^2)}{u} \exp\left(-\frac{u+m_c^2}{T^2}\right) \\
& + \frac{m_s m_c^3 \langle \bar{s}s \rangle}{16\sqrt{2}\pi^2 T^2} \int_{m_c^2}^{s_{D_{s0}}^0} ds \left(1 - \frac{m_c^2}{s}\right)^2 \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& - \frac{m_s m_c \langle \bar{s}s \rangle}{48\sqrt{2}\pi^2 T^2} \int_{m_c^2}^{s_{D_s^*}^0} du \left(1 - \frac{m_c^2}{u}\right)^2 (2u+m_c^2) \exp\left(-\frac{u+m_c^2}{T^2}\right) \\
& - \frac{m_c^3 \langle \bar{s}g_s \sigma G s \rangle}{32\sqrt{2}\pi^2 T^4} \int_{m_c^2}^{s_{D_{s0}}^0} ds \left(1 - \frac{m_c^2}{s}\right) \frac{m_s(s+m_c^2) - m_c(s-m_c^2)}{s} \exp\left(-\frac{s+m_c^2}{T^2}\right) \\
& - \frac{m_c^2 \langle \bar{s}g_s \sigma G s \rangle}{96\sqrt{2}\pi^2 T^4} \int_{m_c^2}^{s_{D_s^*}^0} du \left(1 - \frac{m_c^2}{u}\right) \frac{6m_s m_c u + (2u+m_c^2)(u-m_c^2)}{u} \exp\left(-\frac{u+m_c^2}{T^2}\right)
\end{aligned}$$

$$\begin{aligned}
& + \frac{m_s m_c \langle \bar{s} g_s \sigma G s \rangle}{96 \sqrt{2} \pi^2 T^2} \left(1 + \frac{m_c^2}{T^2} - \frac{m_c^4}{T^4} \right) \int_{m_c^2}^{s_{D_{s0}}^0} ds \left(1 - \frac{m_c^2}{s} \right)^2 \exp \left(-\frac{s+m_c^2}{T^2} \right) + \frac{m_s m_c^3 \langle \bar{s} g_s \sigma G s \rangle}{288 \sqrt{2} \pi^2 T^6} \\
& \times \int_{m_c^2}^{s_{D_s^*}^0} du \left(1 - \frac{m_c^2}{u} \right)^2 (2u + m_c^2) \exp \left(-\frac{u+m_c^2}{T^2} \right) + \frac{m_c^2 \langle \bar{s} g_s \sigma G s \rangle}{96 \sqrt{2} \pi^2} \int_{m_c^2}^{s_{D_{s0}}^0} ds \frac{m_s m_c - 2s + m_c^2}{s^2} \exp \left(-\frac{s+m_c^2}{T^2} \right) \\
& + \frac{m_c \langle \bar{s} g_s \sigma G s \rangle}{96 \sqrt{2} \pi^2} \int_{m_c^2}^{s_{D_s^*}^0} du \frac{2m_s u + m_c^3}{u^2} \exp \left(-\frac{u+m_c^2}{T^2} \right),
\end{aligned} \tag{C33}$$

$$\begin{aligned}
\Pi_{J/\psi \phi S \tilde{V}}^{\text{QCD}}(T^2) &= \frac{9m_s m_c}{256 \sqrt{2} \pi^4} \int_{4m_c^2}^{s_{J/\psi}^0} ds \int_0^{s_{D_s}^0} du \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \exp \left(-\frac{s+u}{T^2} \right) - \frac{3m_c \langle \bar{s} s \rangle}{32 \sqrt{2} \pi^2} \int_{4m_c^2}^{s_{J/\psi}^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \exp \left(-\frac{s}{T^2} \right) \\
&+ \frac{4m_c \langle \bar{s} g_s \sigma G s \rangle}{128 \sqrt{2} \pi^2 T^2} \int_{4m_c^2}^{s_{J/\psi}^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \exp \left(-\frac{s}{T^2} \right) - \frac{5m_c \langle \bar{s} g_s \sigma G s \rangle}{384 \sqrt{2} \pi^2} \int_{4m_c^2}^{s_{J/\psi}^0} ds \frac{1}{\sqrt{s(s-4m_c^2)}} \exp \left(-\frac{s}{T^2} \right),
\end{aligned} \tag{C34}$$

$$\begin{aligned}
\Pi_{D_{s1} D_s S \tilde{V}}^{\text{QCD}}(T^2) &= \frac{1}{64 \sqrt{2} \pi^4} \int_{m_c^2}^{s_{D_{s1}}^0} ds \int_{m_c^2}^{s_{D_s}^0} du \left(1 - \frac{m_c^2}{s} \right) \left(1 - \frac{m_c^2}{u} \right) \exp \left(-\frac{s+u}{T^2} \right) \\
&\frac{m_c (2s + m_c^2) (s - m_c^2) (u - m_c^2) - m_s [m_c^6 + m_c^4 (u - 5s) + m_c^2 s (7u - 2s) - 2s^2 u]}{su} \\
&+ \frac{\langle \bar{s} s \rangle}{24 \sqrt{2} \pi^2} \int_{m_c^2}^{s_{D_{s1}}^0} ds \left(1 - \frac{m_c^2}{s} \right) \frac{6m_s m_c s - (2s + m_c^2) (s - m_c^2)}{s} \exp \left(-\frac{s+m_c^2}{T^2} \right) \\
&+ \frac{m_c \langle \bar{s} s \rangle}{8 \sqrt{2} \pi^2} \int_{m_c^2}^{s_{D_s}^0} du \left(1 - \frac{m_c^2}{u} \right) \frac{m_s (u + m_c^2) + m_c (u - m_c^2)}{u} \exp \left(-\frac{u+m_c^2}{T^2} \right) \\
&+ \frac{m_s m_c \langle \bar{s} s \rangle}{48 \sqrt{2} \pi^2 T^2} \int_{m_c^2}^{s_{D_{s1}}^0} ds \left(1 - \frac{m_c^2}{s} \right)^2 (2s + m_c^2) \exp \left(-\frac{s+m_c^2}{T^2} \right) \\
&- \frac{m_s m_c^3 \langle \bar{s} s \rangle}{16 \sqrt{2} \pi^2 T^2} \int_{m_c^2}^{s_{D_s}^0} du \left(1 - \frac{m_c^2}{u} \right)^2 \exp \left(-\frac{u+m_c^2}{T^2} \right) \\
&- \frac{m_c^2 \langle \bar{s} g_s \sigma G s \rangle}{96 \sqrt{2} \pi^2 T^4} \int_{m_c^2}^{s_{D_{s1}}^0} ds \left(1 - \frac{m_c^2}{s} \right) \frac{6m_s m_c s - (2s + m_c^2) (s - m_c^2)}{s} \exp \left(-\frac{s+m_c^2}{T^2} \right) \\
&- \frac{m_c^3 \langle \bar{s} g_s \sigma G s \rangle}{32 \sqrt{2} \pi^2 T^4} \int_{m_c^2}^{s_{D_s}^0} du \left(1 - \frac{m_c^2}{u} \right) \frac{m_s (u + m_c^2) + m_c (u - m_c^2)}{u} \exp \left(-\frac{u+m_c^2}{T^2} \right) \\
&- \frac{m_s m_c^3 \langle \bar{s} g_s \sigma G s \rangle}{288 \sqrt{2} \pi^2 T^6} \int_{m_c^2}^{s_{D_{s1}}^0} ds \left(1 - \frac{m_c^2}{s} \right)^2 (2s + m_c^2) \exp \left(-\frac{s+m_c^2}{T^2} \right) \\
&- \frac{m_s m_c \langle \bar{s} g_s \sigma G s \rangle}{96 \sqrt{2} \pi^2 T^2} \left(1 + \frac{m_c^2}{T^2} - \frac{m_c^4}{T^4} \right) \int_{m_c^2}^{s_{D_s}^0} du \left(1 - \frac{m_c^2}{u} \right)^2 \exp \left(-\frac{u+m_c^2}{T^2} \right) \\
&+ \frac{m_c \langle \bar{s} g_s \sigma G s \rangle}{96 \sqrt{2} \pi^2} \int_{m_c^2}^{s_{D_{s1}}^0} ds \frac{2m_s s - m_c^3}{s^2} \exp \left(-\frac{s+m_c^2}{T^2} \right) \\
&+ \frac{m_c^2 \langle \bar{s} g_s \sigma G s \rangle}{96 \sqrt{2} \pi^2} \int_{m_c^2}^{s_{D_s}^0} du \frac{m_s m_c + 2u - m_c^2}{u^2} \exp \left(-\frac{u+m_c^2}{T^2} \right),
\end{aligned} \tag{C35}$$

$$\begin{aligned}\Pi_{\eta_c\phi S\tilde{V}}^{\text{QCD}}(T^2) = & \frac{m_c}{16\sqrt{2}\pi^4} \int_{4m_c^2}^{s_{\eta_c}^0} ds \int_0^{s_\phi^0} du u \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \exp\left(-\frac{s+u}{T^2}\right) - \frac{m_s m_c \langle \bar{s}s \rangle}{2\sqrt{2}\pi^2} \int_{m_c^2}^{s_{\eta_c}^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \exp\left(-\frac{s}{T^2}\right) \\ & - \frac{m_s m_c \langle \bar{s}g_s \sigma G s \rangle}{24\sqrt{2}\pi^2} \int_{m_c^2}^{s_{\eta_c}^0} ds \frac{1}{\sqrt{s(s-4m_c^2)}} \exp\left(-\frac{s}{T^2}\right),\end{aligned}\quad (\text{C36})$$

$$\begin{aligned}\Pi_{\chi_{c1}\phi S\tilde{V}}^{\text{QCD}}(T^2) = & \frac{1}{48\sqrt{2}\pi^4} \int_{4m_c^2}^{s_{\chi_{c1}}^0} ds \int_0^{s_\phi^0} du \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} u (s-4m_c^2) \exp\left(-\frac{s+u}{T^2}\right) - \frac{m_s \langle \bar{s}s \rangle}{6\sqrt{2}\pi^2} \int_{4m_c^2}^{s_{\chi_{c1}}^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \\ & \times (s-4m_c^2) \exp\left(-\frac{s}{T^2}\right) + \frac{m_s m_c^2 \langle \bar{s}g_s \sigma G s \rangle}{24\sqrt{2}\pi^2} \int_{4m_c^2}^{s_{\chi_{c1}}^0} ds \frac{1}{\sqrt{s(s-4m_c^2)}} \exp\left(-\frac{s}{T^2}\right),\end{aligned}\quad (\text{C37})$$

$$\begin{aligned}\Pi_{\chi_{c0}\phi S\tilde{V}}^{\text{QCD}}(T^2) = & \frac{3m_s}{32\sqrt{2}\pi^4} \int_{4m_c^2}^{s_{\chi_{c0}}^0} ds \int_0^{s_\phi^0} du \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} (s-4m_c^2) \exp\left(-\frac{s+u}{T^2}\right) \\ & - \frac{\langle \bar{s}s \rangle}{4\sqrt{2}\pi^2} \int_{4m_c^2}^{s_{\chi_{c0}}^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} (s-4m_c^2) \exp\left(-\frac{s}{T^2}\right) \\ & + \frac{\langle \bar{s}g_s \sigma G s \rangle}{12\sqrt{2}\pi^2 T^2} \int_{4m_c^2}^{s_{\chi_{c0}}^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} (s-4m_c^2) \exp\left(-\frac{s}{T^2}\right) \\ & - \frac{\langle \bar{s}g_s \sigma G s \rangle}{24\sqrt{2}\pi^2} \int_{4m_c^2}^{s_{\chi_{c0}}^0} ds \frac{s-5m_c^2}{\sqrt{s(s-4m_c^2)}} \exp\left(-\frac{s}{T^2}\right),\end{aligned}\quad (\text{C38})$$

$$\begin{aligned}\Pi_{J/\psi f_0(980)S\tilde{V}}^{\text{QCD}}(T^2) = & \frac{3m_c}{32\sqrt{2}\pi^4} \int_{4m_c^2}^{s_{J/\psi}^0} ds \int_0^{s_{f_0(980)}^0} du u \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \exp\left(-\frac{s+u}{T^2}\right) \\ & + \frac{m_s m_c \langle \bar{s}s \rangle}{2\sqrt{2}\pi^2} \int_{4m_c^2}^{s_{J/\psi}^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \exp\left(-\frac{s}{T^2}\right) \\ & + \frac{m_s m_c \langle \bar{s}g_s \sigma G s \rangle}{12\sqrt{2}\pi^2} \int_{4m_c^2}^{s_{J/\psi}^0} ds \frac{s-m_c^2}{s\sqrt{s(s-4m_c^2)}} \exp\left(-\frac{s}{T^2}\right) \\ & + \frac{m_s m_c \langle \bar{s}g_s \sigma G s \rangle}{4\sqrt{2}\pi^2 T^2} \int_{4m_c^2}^{s_{J/\psi}^0} ds \frac{\sqrt{\lambda(s, m_c^2, m_c^2)}}{s} \exp\left(-\frac{s}{T^2}\right),\end{aligned}\quad (\text{C39})$$

$$\lambda(a, b, c) = a^2 + b^2 + c^2 - 2ab - 2bc - 2ca.$$

References

- [1] S. Navas *et al.* (Particle Data Group), *Phys. Rev. D* **110**, 030001 (2024)
- [2] X. L. Wang *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **99**, 142002 (2007)
- [3] J. P. Lees *et al.* (BaBar Collaboration), *Phys. Rev. D* **89**, 111103 (2014)
- [4] X. L. Wang *et al.* (Belle Collaboration), *Phys. Rev. D* **91**, 112007 (2015)
- [5] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **130**, 121901 (2023)
- [6] G. Pakhlova *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **101**, 172001 (2008)
- [7] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **131**, 191901 (2023)
- [8] S. Jia *et al.* (Belle Collaboration), *Phys. Rev. D* **100**, 111103 (2019)
- [9] S. Jia *et al.* (Belle Collaboration), *Phys. Rev. D* **101**, 091101 (2020)

- [10] F. K. Guo, J. Haidenbauer, C. Hanhart *et al.*, *Phys. Rev. D* **82**, 094008 (2010)
- [11] G. Cotugno, R. Faccini, A. D. Polosa *et al.*, *Phys. Rev. Lett.* **104**, 132005 (2010)
- [12] D. V. Bugg, *J. Phys. G* **36**, 075002 (2009)
- [13] F. K. Guo, C. Hanhart, and U. G. Meissner, *Phys. Lett. B* **665**, 26 (2008)
- [14] R. M. Albuquerque, M. Nielsen, and R. Rodrigues da Silva, *Phys. Rev. D* **84**, 116004 (2011)
- [15] Z. G. Wang and X. H. Zhang, *Commun. Theor. Phys.* **54**, 323 (2010)
- [16] R. M. Albuquerque and M. Nielsen, *Nucl. Phys. A* **815**, 53 (2009) [Erratum: *Nucl. Phys. A* **857**, 48 (2011)]
- [17] D. Ebert, R. N. Faustov, and V. O. Galkin, *Eur. Phys. J. C* **58**, 399 (2008)
- [18] W. Chen and S. L. Zhu, *Phys. Rev. D* **83**, 034010 (2011)
- [19] J. R. Zhang and M. Q. Huang, *Phys. Rev. D* **83**, 036005 (2011)
- [20] M. N. Anwar, J. Ferretti, and E. Santopinto, *Phys. Rev. D* **98**, 094015 (2018)
- [21] H. Sundu, S. S. Agaev, and K. Azizi, *Phys. Rev. D* **98**, 054021 (2018)
- [22] L. Maiani, F. Piccinini, A. D. Polosa *et al.*, *Phys. Rev. D* **89**, 114010 (2014)
- [23] Z. G. Wang, *Eur. Phys. J. C* **78**, 518 (2018)
- [24] Z. G. Wang, *Eur. Phys. J. C* **74**, 2874 (2014)
- [25] Z. G. Wang, *Eur. Phys. J. C* **76**, 387 (2016)
- [26] Z. G. Wang, *Nucl. Phys. B* **973**, 115592 (2021)
- [27] Z. G. Wang, *Nucl. Phys. B* **1002**, 116514 (2024)
- [28] N. Li, H. Z. He, W. Liang *et al.*, *Chin. Phys. C* **47**, 063102 (2023)
- [29] S. Dubynskiy and M. B. Voloshin, *Phys. Lett. B* **666**, 344 (2008)
- [30] G. J. Ding, J. J. Zhu, and M. L. Yan, *Phys. Rev. D* **77**, 014033 (2008)
- [31] M. N. Anwar, Y. Lu, and B. S. Zou, *Phys. Rev. D* **95**, 114031 (2017)
- [32] Q. Deng, R. H. Ni, Q. Li *et al.*, *Phys. Rev. D* **110**, 056034 (2024)
- [33] L. Y. Xiao, X. Z. Weng, Q. F. Lü *et al.*, *Eur. Phys. J. C* **78**, 605 (2018)
- [34] B. Q. Li and K. T. Chao, *Phys. Rev. D* **79**, 094004 (2009)
- [35] J. Z. Wang, R. Q. Qian, X. Liu *et al.*, *Phys. Rev. D* **101**, 034001 (2020)
- [36] B. D. Wan, L. Tang, and C. F. Qiao, *Eur. Phys. J. C* **80**, 121 (2020)
- [37] C. F. Qiao, *Phys. Lett. B* **639**, 263 (2006)
- [38] C. F. Qiao, *J. Phys. G* **35**, 075008 (2008)
- [39] L. Q. Song, D. Song, J. T. Zhu *et al.*, *Phys. Lett. B* **835**, 137586 (2022)
- [40] Z. Liu, H. Xu, Z. W. Liu *et al.*, arXiv: 2603.09404 [hep-ph]
- [41] X. W. Wang, Z. G. Wang, and G. L. Yu, *Eur. Phys. J. A* **57**, 275 (2021)
- [42] Z. G. Wang, *Front. Phys.* **21**, 016300 (2026)
- [43] Z. G. Wang, *Phys. Rev. D* **101**, 074011 (2020)
- [44] R. M. Albuquerque, J. M. Dias, K. P. Khemchandani *et al.*, *J. Phys. G* **46**, 093002 (2019)
- [45] B. Aubert *et al.* (BaBar Collaboration), *Phys. Rev. Lett.* **95**, 142001 (2005)
- [46] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **96**, 032004 (2017)
- [47] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **104**, 052012 (2021)
- [48] Z. G. Wang, *Chin. Phys. C* **48**, 103103 (2024)
- [49] Z. G. Wang, *Eur. Phys. J. C* **78**, 933 (2018)
- [50] Z. G. Wang, *Eur. Phys. J. C* **79**, 29 (2019)
- [51] Z. G. Wang, *Eur. Phys. J. C* **79**, 184 (2019)
- [52] M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, *Nucl. Phys. B* **147**, 385 (1979); *Nucl. Phys. B* **147**, 448 (1979)
- [53] L. J. Reinders, H. Rubinstein, and S. Yazaki, *Phys. Rept.* **127**, 1 (1985)
- [54] Z. G. Wang and J. X. Zhang, *Eur. Phys. J. C* **78**, 14 (2018)
- [55] Z. G. Wang and X. S. Yang, *AAPPs Bull.* **34**, 5 (2024)
- [56] X. S. Yang and Z. G. Wang, *Chin. Phys. C* **49**, 063108 (2025)
- [57] Z. G. Wang, *Eur. Phys. J. C* **75**, 427 (2015)
- [58] P. Ball and G. W. Jones, *JHEP* **0703**, 069 (2007)
- [59] H. Y. Cheng, C. W. Chiang, and Z. Q. Zhang, *Phys. Rev. D* **105**, 033006 (2022)
- [60] Z. G. Wang, *Eur. Phys. J. C* **76**, 427 (2016)
- [61] Z. G. Wang, W. M. Yang, and S. L. Wan, *Eur. Phys. J. C* **37**, 223 (2004)
- [62] D. Becirevic, G. Duplancic, B. Klajn *et al.*, *Nucl. Phys. B* **883**, 306 (2014)
- [63] V. A. Novikov, L. B. Okun, M. A. Shifman *et al.*, *Phys. Rept.* **41**, 1 (1978)
- [64] P. Colangelo and A. Khodjamirian, arXiv: hep-ph/0010175
- [65] S. Narison and R. Tarrach, *Phys. Lett. B* **125**, 217 (1983)
- [66] Z. G. Wang and T. Huang, *Phys. Rev. D* **89**, 054019 (2014)
- [67] Z. G. Wang and T. Huang, *Eur. Phys. J. C* **74**, 2891 (2014)
- [68] Z. G. Wang, *Eur. Phys. J. C* **74**, 2963 (2014)
- [69] Z. G. Wang and T. Huang, *Nucl. Phys. A* **930**, 63 (2014)
- [70] R. M. Albuquerque, S. Narison, A. Rabemananjara, and D. Rabetiariivony, *Nucl. Part. Phys. Proc.* **343**, 61 (2024)
- [71] R. M. Albuquerque, S. Narison, and D. Rabetiariivony, *Nucl. Part. Phys. Proc.* **324-329**, 54 (2023)
- [72] R. H. Wua, C. Y. Wang, C. Meng *et al.*, *JHEP* **06**, 216 (2024)